APPENDIX C

Preliminary Delineation of Wetlands and Other Waters Report



Guadalupe River Bridge Replacement Project Preliminary Delineation of Wetlands and Other Waters



Prepared for: Peninsula Corridor Joint Powers Board 1250 San Carlos Avenue San Carlos, California 94070

> Prepared by: MIG 2055 Junction Avenue Suite 205 San Jose, CA 95131 (650) 327-0429

> > August 2020

PLANNING | DESIGN | COMMUNICATIONS | MANAGEMENT | SCIENCE | TECHNOLOGY

2055 Junction Avenue, Suite 205 • San Jose, CA 95131 • USA • 650-327-0429 • www.migcom.com Offices in: California • Colorado • Oregon • Texas • Washington

Table of Contents

Executive Summary					
1.	I. Introduction				
	2.1	Project Study Area Description	5		
	2.2	Proposed Project	6		
	2.3	Project Purpose	7		
2.	Surve	y Methods	13		
	2.1	Identification of Jurisdictional Waters	.13		
	2.2	Identification of Section 404 Wetlands (Special Aquatic Sites)	.14		
	2.3	Identification of Section 404 Jurisdictional Other Waters	.16		
	2.4	Identification of Waters of the State	.16		
	2.5	Identification of CDFW Jurisdiction	.17		
3.	Surve	y Results and Discussion	18		
	3.1	Study Area Conditions and Observations	.20		
	3.2	Rationale for Sample Points	.21		
	3.3	Photo Points	.23		
	3.4	Identification of Section 404 Potentially Jurisdictional Waters	.24		
	3.5	Identification of Section 404 Potentially Jurisdictional Wetlands	.24		
	3.6	Identification of Potentially Jurisdictional Waters of the State	.25		
	3.7	Identification of CDFW Potentially Jurisdictional Habitats	.25		
	3.8	Areas Not Meeting the Regulatory Definition of Section 404/401 Wetlands and Waters	.27		
4.	Refere	ences	30		
Арреі	ndix A:	Soil Report for the Study Area	32		
Apper	ndix B:	Plants Observed in the Study Area	33		
Appendix C: USACE Western Mountains, Valley and Coast Wetland Data Forms 36					
Appendix D: Photographic Documentation of the Study Area					
Apper	Appendix E: Aquatic Resources Table43				

List Of Tables

Table 1. Classification of Wetland-Associated Plant Species (Lichvar et al. 2016)15
Table 2. Summary of Jurisdictional Waters and Habitats within the Study Area	
Table 3. Summary of OHWM Sample Points within the Study Area	
Table 4. Coordinates and Rationale for Photo Points	24

List of Figures

Figure 1 Vicinity Map	8
Figure 2 Project Site Map	9
Figure 3 USGS Topographic Map	10
Figure 4 NRCS Soils Map	11
Figure 5 NWI Map	12
Figure 6 Vegetation Communities Map	
Figure 7 Preliminary Identification of Waters of the U.S./State	29

Executive Summary

MIG surveyed the Guadalupe River Bridge Replacement Project study area located in San Jose, Santa Clara County, California for wetlands and other waters potentially subject to regulation under Section 404 of the Clean Water Act (CWA) as administered by the United States Army Corps of Engineers (USACE). The survey also delineated the extent of waters of the state that may be subject to regulation by the Regional Water Quality Control Board (RWQCB) under Section 401 of the Clean Water Act and under the Porter Cologne Water Quality Control Act. Lastly, the survey also delineated jurisdictional habitats subject to regulation under Sections 1600-1607 of the California Fish and Game Code, which is administered by the California Department of Fish and Wildlife (CDFW).

In total, approximately 4.39 acres of potentially USACE and RWQCB jurisdictional features were identified in the study area. These include approximately 1.45 acres of Sections 401 and 404 waters situated below the ordinary high water mark (OHWM) of Guadalupe River. Section 401 and 404 wetlands are also present throughout the study area and total 2.94 acres. Section 401 waters of the state extend farther up to the top of the banks for an additional 1.66 acres.

CDFW jurisdictional features as defined by bed and bank topography, and riparian habitat were identified in the study area and total 6.67 acres, including aquatic, wetlands, riparian habitat, annual grassland habitat, and developed land within top of bank, as well as riparian habitat extending beyond top of bank.

1. Introduction

2.1 Project Study Area Description

Caltrain, a commuter rail line on the San Francisco Peninsula and in the Santa Clara Valley, owns and operates trains on two tracks (MT-1 and MT-2), between San Francisco and Gilroy. Caltrain is governed by the Peninsula Corridor Joint Powers Board (PCJPB) which consists of agencies from the three counties served by Caltrain, including Santa Clara County. In the study area, the tracks span the Guadalupe River on separate and adjacent bridges, also owned by Caltrain. The 17.06-acre study area for the delineation is surrounded by dense residential development and is located just south of downtown San Jose (Figures 1 and 2). The study area is situated in the *San Jose West* U.S. Geological Survey (USGS) 7.5-minute quadrangle (Figure 3). Elevation of the study area is approximately 86 to 110 feet North American Vertical Datum of 1988 NAVD88 (Google Inc. 2019).

The climate at the study area is coastal Mediterranean, with most rain falling in the winter and spring. Cool to mild temperatures are common in the winter and hot to mild temperatures are common in the summer. Climate conditions in the study area include a 30-year average of approximately 14.9 inches of annual precipitation with an average temperature range from 50.6°F to 72.4°F (PRISM Climate Group 2019). Relative to the 30-year climate normal, the study area experienced normal conditions during the 2018 wet season prior to the December 2018 survey. From September 2018 through November 2018, the area received 1.95 inches of precipitation, which is approximately 68% of the 30-year average for this same period (PRISM Climate Group 2019).

Figure 4 shows the one soil unit mapped by the National Resource Conservation Service (NRCS) in the study area: 165 – Urbanland-Campbell complex, 0 to 2 percent slopes, protected (NRCS 2019a). This soil series is listed as hydric in Santa Clara County on the National Hydric Soils List (NRCS 2019b). A detailed description of this soil type is provided in Appendix A.

The U.S. Fish and Wildlife Service's National Wetlands Inventory (NWI) map of the study area is depicted in Figure 5. The NWI identified Guadalupe River within the study area as a temporarily flooded, palustrine forested/shrub wetland (PFOA) (NWI 2019). NWI maps are based on interpretation of aerial photography, limited verification of mapped units, and/or classification of wetland types using the classification system developed by Cowardin et al. (1979). These data are available for general reference purposes and do not necessarily correspond to the presence or absence of jurisdictional waters.

The Federal Emergency Management Agency (FEMA) produces maps depicting flood zones that are generally associated with rivers, oceans, and other water bodies. Like the NWI maps, the FEMA flood zone maps are based predominantly on topography and regional modeling. Based upon a review of the FEMA flood zone maps of the study area vicinity, the study area lies within Special Flood Hazard Areas zones A and AO with an estimated flood depth of 1 foot in a 100-year (1%) flood event (FEMA 2019).

Guadalupe River Bridge Replacement Project Preliminary Delineation of Wetlands and Other Waters August 2020

2.2 Proposed Project

The PCJPB proposes to replace the damaged and aged MT-1 rail bridge over the Guadalupe River and extend the length of the MT-2 bridge on the right bank of the Guadalupe River. The downstream (northerly) bridge consists of a wooden trestle bridge constructed in 1935 (MT-1) and the upstream bridge consists of a concrete bridge constructed in 1990 as part of the Caltrans Highway 87 project (MT-2). In addition to Caltrain's passenger service, the railroad bridges are used by Union Pacific Railroad (UPRR) freight service; Amtrak passenger service; and by the Altamont Commuter Express (ACE) and Capitol Corridor to reach the Tamien Yard.

The 1935 MT-1 bridge urgently needs to be replaced with a new structure in order to maintain safe and reliable operations for all users. The MT-1 bridge does not meet current railroad structural design standards (including seismic criteria), and as a result is vulnerable to collapse in the event of a significant earthquake. The timber structure of MT-1 has been further damaged by multiple fires, most recently a large fire in November 2017.

The MT-1 and MT-2 bridges are located along a sharp meander of the Guadalupe River and the river exhibits a high degree of floodplain fill, channel confinement and bank failures. The geomorphic issues directly affect the safety and reliability of the railroad bridges by eroding directly towards the bridge abutments. River bank failures at MT-2 occurred in 2017 and at both MT-1 and MT-2 in several previous years, requiring emergency bank stabilization measures. To address these safety issues and protect the rail bridge asset, Caltrain proposes to widen the channel, replace the MT-1 bridge with a new, longer bridge, and to extend the MT-2 bridge. The existing MT-2 bridge meets seismic criteria and does not require replacement but will be lengthened on the south side to help address geomorphic stability issues at the bridge abutments.

The USACE and Santa Clara Valley Water District (Valley Water) propose to implement a separate and independent flood control project in the future, referred to as the Upper Guadalupe River Flood Protection Project (UGRFPP), that includes construction of a bypass channel widening through the project area. While the UGRFPP will not be constructed in the near term, project design has incorporated several measures that do not preclude potential future channel widening and bridge extension.

The existing 187-foot MT-1 bridge will be replaced by a 265-foot pre-cast concrete structure. The center span over the main channel will be 110 feet in length and the pier placement has been optimized through hydraulic analysis to avoid pier placement in the low-flow channel. The bridge piers will consist of two 48-inch diameter cast-in-drilled-hole (CIDH) piles. The new MT-1 bridge will continue to accommodate a single track. Channel widening will occur under the south side of the MT-1 bridge to reduce scour/increase flow capacity. The southern abutment will be designed so that it can potentially function as a pier without modification in the future if the USACE bypass channel is constructed.

The existing 199-foot MT-2 bridge will be extended by 90 feet at the southern end, resulting in a new total bridge length of 244.5 feet. In order to accommodate this extension, the existing MT-2 abutment 5 would be removed and replaced by a new pier and the channel widened. The existing northern abutment 1 and piers 2, 3, and 4 would remain in place. Similar to the MT-1

bridge, the southernmost abutment would be designed to also function as pier if the USACE bypass channel is constructed.

The project also includes the relocation of an existing Overhead Catenary System (OCS) Pole constructed as part of the Peninsula Corridor Electrification Program near the southern abutment of MT-2 and existing fiber optic lines on the MT-1 bridge.

2.3 Project Purpose

The purpose of the project is to address the structural deficiencies of the MT-1 bridge and the geomorphic instability of the Guadalupe River channel in the vicinity of the MT-1 and MT-2 bridges to provide for long- term public safety and service reliability.

Without the project, the structural condition of the MT-1 bridge presents an increasing safety hazard to all users. Replacing MT-1 is needed to meet the standards of safety and reliability required for current and future train loads, to ensure that the bridge will continue to safely carry passenger commuter (Caltrain and Amtrak) and freight (UPRR) rail service well into the future (the bridges are also used for deadhead movements of ACE and Capitol Corridor trains). In addition, lengthening of both the MT-1 and MT-2 bridges is needed to address erosion and scour issues which continue to undermine bridge abutments and contribute to risk of bridge structure failure. Extending both bridges will allow for reduced river flow velocities and minimize bank erosion. Without the project, the geomorphic condition of this reach of the Guadalupe River will continue to contribute to bank failure, threatening the integrity of the transportation asset and requiring continual emergency repair interventions.



Figure 1 Vicinity Map

Guadalupe River Bridge Replacement Project





Study Area (17.06 acres) Guadalupe River

0 80 160 320

Figure 2 Project Site Map Guadalupe River Bridge Replacement Project

MIG



Study Area (17.06 acres)

0 465 930 1,860

Figure 3 USGS Topographic Map Guadalupe River Bridge Replacement Project







Guadalupe River

165 (Urbanland-Campbell complex, 0 to 2 percent slopes, protected)

Figure 4 NRCS Soils Map Guadalupe River Bridge Replacement Project

0

MIG





Study Area (17.06 acres)



MIG

Freshwater Forested/Shrub Wetland (Palustrine, Forested, Temporary Flooded)

340 Feet 170 85

Figure 5 NWI Map

Guadalupe River Bridge Replacement Project

0

2. Survey Methods

Before the delineation survey was conducted, topographic maps and aerial photos of the study area were obtained and reviewed from several sources, such as the USGS (Figure 3), NRCS (Figure 4), NWI (Figure 5), and Google Earth software (Google Inc. 2019).

On December 12, 2018, MIG senior biologist Laura Moran, B.S. and MIG ecologist Charlotte Moran, B.A. performed a technical delineation of wetlands and other waters in the study area, in accordance with the *Corps of Engineers 1987 Wetlands Delineation Manual* (Corps Manual; Environmental Laboratory 1987). Additionally, the *Regional Supplement to the Corps of Engineers Wetland Delineation Manual: Arid West (Version 2.0)* (Regional Supplement) (USACE 2008a) and *A Field Guide to the Identification of the Ordinary High-Water Mark (OHWM) in the Arid West Region of the Western United States* (USACE 2008b) were followed to document site conditions relative to hydrophytic vegetation, hydric soils, and wetland hydrology. The extent and distribution of wetlands and other waters of the U.S. were mapped. These include wetlands and waters that may be subject to regulation under Section 404 of the CWA, and waters of the state that may be subject to regulation under Section 401 of the CWA or the Porter Cologne Water Quality Control Act, which is administered by the RWQCB. The scientists also surveyed for aquatic and riparian habitat that may be subject to regulation under Section 404 of the California Fish and Game Code, which is administered by the California Department of Fish and Wildlife (CDFW).

On August 14, 2020, MIG Senior Biologist David Gallagher, M.S. and MIG biologist Alex Broskoff, B.S. visited the project site to obtain an update on site conditions. This report reflects site conditions observed in August 2020.

2.1 Identification of Jurisdictional Waters

The vegetation, soils, and hydrology in the study area were mapped according to the Routine Determination Method outlined in the Corps Manual (Environmental Laboratory 1987), using updated data forms, vegetation sampling methods, and hydric soil and hydrology indicators developed for the Regional Supplement (USACE 2010). This three-parameter approach to identifying wetlands is based on the presence of a prevalence or dominance of hydrophytic vegetation, hydric soils, and wetland hydrology.

This report was prepared in accordance with guidance provided in *Updated Map and Drawing Standards for the South Pacific Division Regulatory Program* (USACE 2016a) and *Information Requested for Verification of Corps Jurisdiction* (USACE 2016b). These documents list the information that must be submitted as part of a request for a jurisdictional determination, including:

- Vicinity map (Figure 1)
- Project area map (Figure 2)
- USGS quadrangle sheet (Figure 3)
- Soils map (Figure 4)

Guadalupe River Bridge Replacement Project Preliminary Delineation of Wetlands and Other Waters August 2020

- National Wetlands Inventory map (Figure 5)
- Vegetation communities map (Figure 6)
- Delineation map (Figure 7)
- Current soil survey report (Appendix A)
- Plant species observed (Appendix B)
- Arid West Wetland Determination Data Forms (Appendix C)
- Written rationale for sample point choice (Section 3.1, "Observations, Rationales, and
- Assumptions")
- Color photos (Appendix D)
- Aquatic resources table (Appendix E)

During the survey, the study area was examined for topographic features, drainages, alterations to hydrology or vegetation, and recent significant disturbance. A determination was then made as to whether normal environmental conditions were present at the time of the field survey. In the field, the techniques used to identify wetlands included observing the vegetation growing near the soil sample points and characterizing the current surface and subsurface hydrologic features present near the sample points through both observation of indicators and direct observation of hydrology. Features meeting wetland vegetation, soil, and hydrology criteria were then mapped in the field. Geospatial data were collected using a Trimble GeoXT geographic positioning system (GPS) unit with sub-meter accuracy during the 2018 site visit. During the 2020 site visit, geospatial data were collected using a tablet with an Arrow 100 submeter GPS receiver and a geo-spatial mobile-device application.

2.2 Identification of Section 404 Wetlands (Special Aquatic Sites)

Vegetation, soils, and hydrology parameters were recorded where wetland field characteristics were present using the Routine Determination Method outlined in the Corps Manual (Environmental Laboratory 1987) and the updated data forms, vegetation sampling methods, and hydric soil and hydrology indicators developed for the Regional Supplement (USACE 2010).

Hydrophytic Vegetation. Plants that can grow in soils that are saturated or inundated for long periods of time and contain little or no oxygen when wetted, are considered adapted to those soils, and are called hydrophytic. There are different levels of adaptation, as summarized in Table 2. Some plants can only grow in soils saturated with water (and depleted of oxygen), some are mostly found in this condition, and some are found equally in wet soils and in dry soils. Plants observed at each of the sample study areas were identified to species, where possible, using The Jepson Manual, Vascular Plans of California, Second Edition (Baldwin et al. 2012). The wetland indicator status of each species was obtained from the Arid West 2016 Regional Wetland Plant List (Lichvar et al. 2016). Wetland indicator species are designated according to their frequency of occurrence in wetlands. For instance, a species with a presumed frequency of occurrence of 67 to 99 percent in wetlands is designated a facultative wetland indicator species. The wetland indicator groups, indicator symbol, and the frequency of occurrence of species, provided as a percentage, within wetlands are shown in Table 1.

	Indicator Category	Symbol	Frequency (Percent) of Occurrence in Wetlands ¹	
Obligate OBL		OBL	>99 (Almost always is a hydrophyte, rarely in uplands)	
	Facultative wetland	FACW	67 - 99 (Usually a hydrophyte but occasionally found in uplands)	
	Facultative	FAC	34 – 66 (Commonly occurs as either a hydrophyte or non-hydrophyte)	
	Facultative upland	FACU	1 - 33 (Occasionally is a hydrophyte, but usually occurs in uplands)	
	Upland ²	UPL	<1% (Rarely is a hydrophyte, almost always in uplands)	
	Not listed ²	NI	Considered to be an upland species unless otherwise noted	

Table 1.	Classification of	Wetland-Associa	ated Plant Spe	ecies (Lichvar	et al. 2016)
	olassinoation of		aloa i lant ope	Cics (Licitvai	Ct ul. 2010)

Obligate and facultative wetland indicator species are hydrophytes that occur "in areas where the frequency and duration of inundation or soil saturation produce permanently or periodically saturated soils of sufficient duration to exert a controlling influence on the plant species present" (Environmental Laboratory 1987). Facultative indicator species may be considered wetland indicators when found growing in hydric soils that experience periodic saturation. Plant species that are not on the regional list of wetland indicator species are considered upland species. A complete list of the vascular plants observed in the project study area, including their current indicator statuses, is provided in Appendix B.

Hydric Soils. Up to 18 inches of the soil profile were examined for hydric soil indicators. The National Technical Committee for Hydric Soils (NTCHS) defines a hydric soil as one formed under conditions of saturation, flooding or ponding long enough during the growing season to develop anaerobic conditions in the upper 12 inches of soil (NRCS 2010). Hydric soils include soils developed under sufficiently wet conditions to support the growth and regeneration of hydrophytic vegetation. In general, evidence of a hydric soil includes characteristics such as organic soils (histosols), reducing soil conditions, gleyed soils, soils with bright mottles and/or low matrix chroma, soils listed as hydric by the U.S. Department of Agriculture (USDA) on the National Hydric Soils List (NRCS 2019b), and iron and manganese concretions. Reducing soil conditions can also include circumstances where there is evidence of frequent ponding for long or very long duration. A long duration is defined as a period of inundation for a single event that ranges from 7 days to a month and very long is greater than one month (Environmental Laboratory 1987). Munsell Soil Notations were recorded for the soil matrix of each soil sample (Munsell 2009). The Munsell color system is based on three color properties: hue, value, and chroma.

The NRCS Web Soil Survey (NRCS 2019a) was consulted to determine which soil types have been mapped in the project study area (Figure 4). Detailed descriptions of these soil types are provided in Appendix A.

¹ Based on information contained in the Corps Manual.

² Plant species that are not listed in the Arid West 2016 Regional Wetland Plant List (Lichvar et al. 2016) are considered UPL species

Guadalupe River Bridge Replacement Project Preliminary Delineation of Wetlands and Other Waters August 2020

Wetland Hydrology. Wetland hydrology is defined as an area that is inundated either permanently or periodically at mean water depths less than 6.6 feet, or where the soil is saturated at the surface at some time during the growing season of the prevalent vegetation. The period of inundation or soil saturation varies according to the hydrologic/soil moisture regime and occurs in both tidal and non-tidal situations.

Wetland hydrology encompasses all hydrologic characteristics of areas that are periodically inundated or have soils saturated to the surface at some time during the growing season. Wetland hydrology indicators provide evidence that the study area has a continuing wetland hydrologic regime. Primary indicators might include visual observation of surface water (A1), high water table (A2), soil saturation (B1), water-stained leaves (B9), and hydrogen sulfide odor (C1). Secondary indicators might include riverine drift deposits (B3), drainage patterns (B10), and a passing score for the FAC-neutral test (D5). Each of the sample points was examined for positive field indicators (primary and secondary) of wetland hydrology, following the guidance provided in the Regional Supplement.

Potential Section 404 wetlands were identified in the study area.

2.3 Identification of Section 404 Jurisdictional Other Waters

"Other waters" includes lakes, slough channels, seasonal ponds, tributary waters, non-wetland linear drainages, and salt ponds. Such areas are identified by the (seasonal or perennial) presence of standing or running water and generally lack hydrophytic vegetation. In non-tidal or muted tidal waters USACE jurisdiction extends to the ordinary high water mark (OHWM) which is defined in 33 CFR Part 328.3 as "the line on the shore established by the fluctuations of water and indicated by physical characteristics, such as a clear, natural line impressed on the bank, shelving, changes in the character of the soil, destruction of terrestrial vegetation or the presence of litter and debris."

Potential Section 404 other waters were identified in the study area.

2.4 Identification of Waters of the State

The Porter-Cologne Water Quality Control Act (PWQCA) broadly defines waters of the state as "any surface water or groundwater, including saline waters, within the boundaries of the state." Because PWQCA applies to any water, whereas the CWA applies only to certain waters, California's jurisdictional reach overlaps and may exceed the boundaries of waters of the U.S. For example, Water Quality Order No. 2004-0004-DWQ states that "shallow" waters of the state include headwaters, wetlands, and riparian areas. Where forested habitat occurs, the outer canopy of any riparian trees rooted within top of bank may be considered jurisdictional as these trees can provide nutrients and carbon (allochthonous) input to the channel below.

Potential waters of the state were identified in the study area.

Guadalupe River Bridge Replacement Project Preliminary Delineation of Wetlands and Other Waters August 2020

2.5 Identification of CDFW Jurisdiction

Ephemeral and intermittent streams, rivers, creeks, dry washes, sloughs, blue line streams on USGS maps, and watercourses with subsurface flows fall under CDFW jurisdiction. Canals, aqueducts, irrigation ditches, and other means of water conveyance may also be considered streams if they support aquatic life, riparian vegetation, or stream-dependent terrestrial wildlife. A stream is defined in Title 14, California Code of Regulations §1.72, as "a body of water that follows at least periodically or intermittently through a bed or channel having banks and that supports fish and other aquatic life. Jurisdiction does not include tidal areas such as tidal sloughs unless there is freshwater input. This includes watercourses having surface or subsurface flow that supports or has supported riparian vegetation." Using this definition, CDFW extends its jurisdiction to encompass riparian habitats that function as a part of a watercourse. California Fish and Game Code §2786 defines riparian habitat as "lands which contain habitat which grows close to and which depends upon soil moisture from a nearby freshwater source."

The lateral extent of a stream and associated riparian habitat that would fall under the jurisdiction of CDFW can be measured in several ways, depending on the situation and the type of fish or wildlife at risk. At a minimum, CDFW would claim jurisdiction over a stream's bed and bank. For this delineation, the outer edge (dripline) of riparian vegetation was used as the line of demarcation between riparian and upland habitats.

Potential CDFW jurisdictional habitats were identified in the study area.

3. Survey Results and Discussion

The following vegetation/land use communities were mapped in the study area: (1) developed, (2) Wild Oats and Annual Brome Grassland, (3) Fremont Cottonwood Forest, (4) perennial freshwater marsh, (5) seasonal wetland, (6) ornamental woodland (7) aquatic habitat, and (8) Coast Live Oak Woodland (Figure 6).

The parcel is located within the San Francisco Bay Area Subregion of the Central Western Californian Region, both of which are contained within the larger California Floristic Province (Baldwin et al. 2012). Vegetation communities were mapped according to the CDFW Vegetation Classification and Mapping Program's (VegCAMP) currently accepted list of vegetation alliances and associations (CDFW 2020), where applicable.

A total of 24 sample points were examined to identify jurisdictional features (WL 01N/S to WL 12N/S, Appendix C; Figure 7). An additional four sample points were examined during the August 2020 site visit to update jurisdictional features within the study area (SP1 to SP4, Appendix C).

In total, approximately 4.39 acres of potentially USACE and RWQCB jurisdictional features were identified in the study area. These include approximately 1.45 acres of Sections 401 and 404 waters situated below the OHWM of Guadalupe River. Section 401 and 404 wetlands are also present throughout the study area, and total 2.94 acres. Section 401 waters of the state extend farther up to the top of the banks for an additional 1.66 acres. CDFW jurisdictional features as defined by bed and bank topography and riparian habitat were identified in the study area and total 6.67 acres, including aquatic, wetlands, riparian habitat, annual grassland habitat, and developed land within top of bank, as well as riparian habitat extending beyond top of bank. A summary of jurisdictional waters and habitats within the study area is provided in Table 2.

Table 2.	Summary of	Jurisdictional	Waters and	Habitats wit	hin the Study	Area

Potentially Jurisdictional Waters and Habitats	Acres ¹
USACE Jurisdictional Total	4.39
Section 404 Other Waters	
Aquatic habitat	0.73
Fremont Cottonwood Forest (up to OHWM of the Guadalupe River)	0.72
Section 404 Wetlands	
Perennial Freshwater Marsh	1.61
Seasonal Wetland	1.33
RWQCB Jurisdictional Total	6.05
Aquatic Habitat	0.73
Fremont Cottonwood Forest (up to OHWM of the Guadalupe River)	0.72
Fremont Cottonwood Forest (up to the top of bank along the Guadalupe River)	0.79
Perennial Freshwater Marsh	1.61
Seasonal Wetland	1.33
Wild Oats and Annual Brome Grassland (up to top of bank of the Guadalupe River and the flood control basin)	0.67
Developed (up to the top of bank of the Guadalupe River and the flood control basin)	0.20
CDFW Jurisdictional Total	6.67
Aquatic Habitat	0.73
Fremont Cottonwood Forest	2.13
Perennial Freshwater Marsh	1.61
Seasonal Wetland	1.33
Wild Oats and Annual Brome Grassland (up to top of bank of the Guadalupe River and the flood control basin)	0.67
Developed (up to the top of bank of the Guadalupe River and the flood control basin)	0.20

¹Note: Values are approximate due to rounding

Information assembled during this investigation and pertinent to the identification of jurisdictional wetlands and other waters is presented in the five appendices of this report:

- Appendix A—Soil Reports for the Study Area
- Appendix B—Plants Observed in the Study Area
- Appendix C—USACE Western Mountains, Valley and Coast Wetland Data Forms
- Appendix D—Photographic Documentation of the Study Area

Guadalupe River Bridge Replacement Project Preliminary Delineation of Wetlands and Other Waters August 2020

• Appendix E—Aquatic Resources Table

3.1 Study Area Conditions and Observations

- The survey took place during the 2019 wet season and 2020 dry season. Seasonal conditions were considered when assessing the biotic habitats present in the study area. Also, during the 2020 site visit, normal circumstances were not present in the flood control basin due to evidence of recent mowing. However, the boundaries of waters remained clear owing to the presence of hydrology indicators and hydrophytic vegetation.
- A portion of the study area is included Reach 7 of the Upper Guadalupe River Construction Project, a flood damage reduction and recreation project located in the City of San Jose, Santa Clara County, California developed by the Santa Clara Valley Water District (Valley Water) and the USACE. This project begins at Interstate Highway 280, at the edge of downtown San Jose, and extends upstream for approximately 5.5 miles. The proposed project utilizes a combination of bypass channels, floodwalls, and channel widening to achieve flood damage reduction while restoring protected salmonid species habitat (https://www.spn.usace.army.mil/Missions/Projects-and-Programs/Projects-by-Category/Projects-for-Flood-Risk-Management/Upper-Guadalupe-River/).
- The study area is within the San Francisco Bay Sub Region (18050004) of the California Water Resources Region hydrologic unit (USGS 2019).
- Both a seasonal and perennial wetland were mapped within the flood control basin east of Guadalupe River (Figure 6). Based on historic aerial imagery from as early as 1939, the area that contains the flood control basin was within the active floodplain of the river. Based on aerial imagery, the flood control basin was likely constructed in 2010 or 2011. Portions of the basin are regularly inundated and were completely inundated during the winter and spring months of 2017. Based on our site visit in 2020, portions of the basin had been mowed. Also, based on aerial imagery, a portion of the site was possibly graded and filled in late 2019 (Google Inc 2019, 2020; UCSB 2020).
- Along the upper slopes of the banks of the flood control basin, the vegetation is dominated by upland non-native forbs and grasses. This upland vegetation is characterized by wild oat (*Avena fatua*), field bindweed (*Convolvulus arvensis*), and smilo grass (*Stipa miliacea*). Much of the bank slopes are covered with thatch from the last mowing operation.
- Though not relevant to the delineation of waters of the U.S., the top of the bank is mapped for clarity and shown on Figure 7 as Section 401 waters of the State. The current practice of the San Francisco RWQCB is to claim all areas up to the top of bank, but it may also claim riparian habitat that extends beyond the top of bank.

3.2 Rationale for Sample Points

OHWM transects (WL 01N/S to WL 12N/S) and wetland sample points (SP1 to SP4) were selected to document conditions in representative jurisdictional and non-jurisdictional areas (Figure 7).

WL 01N/S to WL 12N/S were selected to demarcate the OHWM of Guadalupe River within the study area. At the time of the delineation, water was flowing and was up to three feet deep. Within the study area, the gradient of the Guadalupe River is low. There are large sections of the bank dominated by rip rap at the bridge crossings as well as downstream and upstream of the bridges. The lower banks were characterized by patchy areas of hydrophytic vegetation, or rip rap, or areas of moderate to severe bank erosion. Generally, the upper banks support a late successional riparian forest. Also, there were several areas along the banks impacted by homeless encampments. Geomorphic field indicators of the OHWM included exposed root hairs and roots below an intact soil layer, break in bank slope, benches formed by differential erosion by change in bank slope, clear, natural line impressed on the bank, and drift (organic and non-organic debris). Vegetative field indicators of the OHWM included vegetation stripped from active areas of the channel, vegetation below OHWM that starts to thicken above OHWM due to lack of disturbance from moderate events, and areas above the OHWM fully vegetated due to lack of disturbance by moderate events. Detailed findings for the OHWM sample point locations are summarized in Table 3.

Table 3. Summary of OHWM Sample Points within the Study Area

Name	Geomorphic Indicator	Vegetative Indicator	Notes
WL 01N/S	Break in bank slope; drift (organic debris)	Vegetation thickens above OHWM due to lack of disturbance from moderate events	Mature riparian forest on upper banks
WL 02 N/S	Benches formed by differential erosion by change in bank slope; drift (organic debris)	Above OHWM is a thick shrub zone indicating lack of physical removal from higher discharge events	Area impacted by homeless encampment; mature riparian forest on upper banks
WL 03 N/S	Break in bank slope	Area above OHWM fully vegetated due to lack of disturbance by moderate events	Mature riparian forest on upper banks
WL 04 N/S	Break in bank slope; benches formed by differential erosion by change in bank slope	Vegetation below OHWM and starts to thicken above OHWW due to lack of disturbance from moderate events	Mature riparian forest on upper banks
WL 05 N/S	Break in bank slope	Vegetation below OHWM and starts to thicken above OHWM due to lack of disturbance from moderate events	Mature riparian forest on upper banks
WL 06 N/S	Break in bank slope	Area above OHWM fully vegetated due to lack of disturbance by moderate events	Young trees rooted within the OHWM; mature riparian forest on upper banks
WL 07 N/S	Benches formed by differential erosion by change in bank slope	Sparse to intermittent vegetation at the edges of the active channel	Mature trees rooted below the OHWM; mature riparian forest on upper banks
WL 08 N/S	Drift deposits (organic and non-organic debris)	Area above OHWM fully vegetated due to lack of disturbance by moderate events	Area impacted by homeless encampment
WL 09 N/S	Benches formed by differential erosion by change in bank slope; drift (organic debris)	Vegetation stripped from active areas of the channel	Area impacted by homeless encampment
WL 10 N/S	Clear line impressed on bent of bridge; exposed roots below intact soil layer; benches formed by differential erosion by change in bank slope	Sparse to intermittent vegetation at the edges of the active channel	Rip rap present; severe bank erosion
WL 11 N/S	Drift (organic debris); break in bank slope	Above OHWM is a thick shrub zone indicating lack of physical removal from higher discharge events	Rip rap present; severe bank erosion
WL 12 N/S	Drift (organic and non-organic debris)	Vegetation below OHWM and starts to thicken above OHWM due to lack of disturbance from moderate events	Rip rap present; severe bank erosion

Guadalupe River Bridge Replacement Project Preliminary Delineation of Wetlands and Other Waters August 2020

SP1 was selected to examine the seasonal wetland in the southern portion of the flood control basin in the study area (Figure 7). Vegetation present was dominated by a single FAC species, bristly ox-tongue (*Helminthotheca echioides*) and the soil exhibited redox depressions. Hydrological indicators, such as saturation visible on aerial imagery and drainage patterns were also observed (Google Inc 2020). The soil was slightly moist. Bristly ox-tongue exhibited increased abundance and plant vigor at SP1, indicating wetland conditions. This is particularly noticeable later in the growing season when adjacent areas are drying out, but moist soils are still present in wetlands, as was the case at SP1

SP2 was chosen to examine the edge of the perennial marsh within the flood control basin near culvert C1 in the study area (Figure 7). Vegetation was dominated by OBL (broadleaf cattail, *Typha latifolia*) and FACW (arroyo willow, *Salix lasiolepis*) species and the soil exhibited a depleted matrix. Hydrological indicators, such as saturation visible on aerial imagery and drainage patterns were also observed (Google Inc 2020). Standing water was observed within 2 feet of SP2 where the perennial marsh continues into a depression. The soil was very moist, but not saturated.

SP3 was selected to investigate the perennial marsh at the northern end of the flood control basin in the study area and was located within a depression (Figure 7.) Vegetation was dominated by a single OBL species, water primrose (*Ludwigia peploides*), and the soil exhibited a depleted matrix. Hydrological indicators, such as saturation visible on aerial imagery and drainage patterns were also observed (Google Inc 2020). The soil was almost saturated, likely indicating the area was recently inundated.

SP4 was chosen to represent uplands along the banks of the flood control basin in the study area (Figure 7). It is located near SP3 in an area sparsely vegetated and dominated by a single upland forb (field bindweed, *Convolvulus arvensis*), and the area was likely mowed recently, based on the presence of a thick layer of thatch. The soil exhibited relic hydric indicators as defined by sharp boundaries between redox concentrations and the soil matrix. Relic indicators are an indication that the soil was likely excavated from the flood plain adjacent to Guadalupe River and used to create the banks of the flood control basin.

3.3 Photo Points

Photo point labels, coordinates, and rationale for the photos are include in Table 4. Photos are included in Appendix D and photo points in Figure 7.

Label	Latitude	Longitude	Rationale
Photo 1	37.316543°	-121.888070°	Guadalupe River and Bridges
Photo 2	37.319272°	-121.888687°	Flood Control Basin
Photo 3	37.318115°	-121.887861°	Seasonal Wetland
Photo 4	37.317576	-121.887634	Perennial Freshwater Marsh
Photo 5	37.317458	-121.887618	Annual Grassland Habitat along Banks of the Flood Control Basin
Photo 6	37.317972	-121.887705	Developed Land Use
Photo 7	37.317898°	-122.888518°	Riparian and Aquatic Habitat

Table 4. Coordinates and Rationale for Photo Points

3.4 Identification of Section 404 Potentially Jurisdictional Waters

Approximately 0.73 acres of Section 404 other waters (open water habitat) were mapped in the study area (Figure 7) and 0.72 acres of Fremont Cottonwood Forest (PS-1) (see Section 3.7 below for a description of this vegetation community). This habitat includes the open water channel (aquatic habitat) and trees rooted below the OHWM of the Guadalupe River (PS1, Figure 7; WL 01N/S to WL 12N/S, Appendix C; Photo 1, Appendix D).

Guadalupe River. The Guadalupe River is an urban, northward flowing 14-mile perennial river that receives flows from creeks that originate in the Santa Cruz Mountains and empties into San Francisco Bay at Alviso Slough. The river begins on the Santa Clara Valley floor at the confluence of Los Alamitos and Guadalupe Creeks. Other major tributaries include Ross Creek, Los Gatos Creek, and Canoas Creek. Guadalupe River is within the Guadalupe watershed, which drains 170 square miles within Santa Clara County. The primary sources of hydrology include a combination of groundwater and seasonal precipitation. The Guadalupe River Basin has been greatly affected by human activity including the urbanization of the surrounding areas, installation of dams and reservoirs, the channelization of streams, and construction of levees for flood protection.

3.5 Identification of Section 404 Potentially Jurisdictional Wetlands

In general, areas that were identified as wetlands included solid stands of hydrophytes and/or areas determined to be ponded and/or saturated for long duration. Approximately 2.94 acres of potential USACE jurisdictional wetlands were identified in the study area (Figure 7). These areas included seasonal wetlands (SW-1 and SW-2) and a perennial freshwater marsh (PM-1). Three parameters identifying Section 404 wetlands were observed at three sample points (Figure 7; SP1 to SP3, Appendix C). The features that were determined to be potentially USACE jurisdictional wetlands are summarized below.

Seasonal Wetland (SW-1 and SW-2). Seasonal wetlands are generally inundated by shallow water, or have high groundwater levels, for variable periods from winter to spring, but they may

be completely dry for most of the summer and fall. Dominant vegetation can include strongly hydrophytic vegetation when the wetland is inundated or saturated and non-hydrophytic, upland species after the wetland dries out. Approximately 1.33 acres of seasonal wetland was mapped within the flood control basin within the study area (SW1 and SW2, Figure 7; SP1, Appendix C; Photo 3, Appendix D). The seasonal wetlands extend from the edge of the perennial marsh up to the toe of the flood control basin banks. The main source of hydrology is likely from a high groundwater table. Dominant species observed included bristly ox-tongue, bird's foot trefoil (*Lotus corniculatus*, FAC), California mugwort (*Artemisia douglasiana*, FAC), and tall flatsedge (*Cyperus eragrostis*, FACW).

Perennial Freshwater Marsh (PM-1). Perennial marshes are generally inundated or have high groundwater levels year-round or for extended periods, but surface water may be lacking during the summer and fall. Approximately 1.61 acres of perennial marsh was mapped within the flood control basin (PM1, Figure 7; SP2, Appendix C; Photo 4, Appendix D). The perennial marsh was mostly confined to a network of depressions within the flood control basin. The source of hydrology is surface flow from culvert C1 and likely from a high groundwater table. Standing water, up to 1 foot deep was observed at C1. Dominant species observed included broadleaf cattail, arroyo willow, water primrose, and rough cocklebur (*Xanthium strumarium*, FACW).

3.6 Identification of Potentially Jurisdictional Waters of the State

The extent of Section 401 waters of the state (RWQCB jurisdiction) in the study area includes a total of 6.05 acres, including areas within Section 404 jurisdiction as described above and riparian habitat, grassland habitat, and developed land up to the top of the banks. In the field, the top of bank was determined by mapping the first significant topographic break in slope. Waters of the state within the study area include all waters of the U.S., and cover approximately, 1.33 acres of seasonal wetland, 0.73 acres of aquatic habitat, 1.61 acres of perennial marsh, 1.51 acres of Fremont Cottonwood Forest up to the top of bank, 0.67 acres of Wild Oats and Annual Brome Grassland, and 0.20 acres of developed land up to the top of bank (Figure 7). Characteristics of waters of the U.S., including wetlands are described above in Sections 3.4 and 3.5. Habitats mapped along the banks are described in Section 3.7 below.

3.7 Identification of CDFW Potentially Jurisdictional Habitats

The study area contains a perennial stream channel with defined bed and bank topography along with associated riparian habitat, as defined by CDFW. Riparian habitat was mapped by the dripline of trees at the outer extent of riparian vegetation. Streambed features were mapped by the top of bank (which can extend beyond the OHWM that is used to measure the extent of waters of the U.S.). The top of bank was delineated in the field as the first distinct topographic break in bank slope. Approximately 0.73 acres of aquatic habitat (up to the OHWM), and 1.51 acres of Fremont Cottonwood Forest up to the top of bank of the streambed plus 0.62 acres of Fremont Cottonwood Forest that extends beyond the top of bank of the streambed, 0.67 acres of Wild Oats and Annual Brome Grassland, 0.20 acres of developed areas along the bank slopes of the Guadalupe River and the flood control basin, 1.33 acres of seasonal wetland, and

1.61 acres of perennial wetland within the banks of the flood control basin are identified as potentially within CDFW jurisdiction.

Fremont Cottonwood Forest (*Populus fremontii* – Salix *laevigata*, S. *lasiolepis* Alliance). Approximately 2.28 acres of this riparian community extends from the water's edge and up the bank of Guadalupe River. A portion of this community was mapped below the OHWM of the river and was included as waters of the U.S. These areas mainly consisted of deeply rooted trees and generally lacked an herbaceous understory or emergent vegetation, likely due to being within the active part of the channel. The large sections of the banks dominated by rip rap at the bridge crossings as well as downstream and upstream of the bridges were mapped as developed land (see below). The dominant trees included Fremont cottonwood and red willow (*Salix laevigata*) with lesser numbers of boxelder (*Acer negundo*). Within the study area, the canopy is intermediate to continuous. Dominant shrubs observed consisted of arroyo willow. Species observed in the open to dense understory above the OHWM included sweet fennel (*Foeniculum vulgare*), curly dock (*Rumex crispus*), broadleaf cattail, smilo grass, white horehound (*Marrubium vulgare*), Himalayan blackberry (*Rubus armeniacus*), English ivy (*Hedera helix*), tall flatsedge, poison hemlock (*Conium maculatum*), and giant reed (*Arundo donax*).

Wild Oats and Annual Brome Grassland. This annual grassland habitat is dominated by nonnative grasses, including wild oats (*Avena* sp.), ripgut brome (*Bromus diandrus*), Mediterranean barley (*Hordeum murinum*), and smilo grass. Other species observed included non-native plant species that are characteristic of disturbed areas, including black mustard (*Brassica nigra*), wild radish (*Raphanus sativus*), Canada horseweed (*Erigeron canadensis*), field bindweed, Italian thistle (*Carduus pycnocephalus*), prickly lettuce (*Lactuca serriola*), and Russian thistle (*Salsola tragus*). Annual grassland was mapped within the top of bank of the Guadalupe River and the flood control basin as well as areas outside of the banks of the river and basin.

Developed Land. Developed land within the top of bank included areas that are covered in rip rap or hardpacked soil that does not support vegetation. These areas were mapped under the bridge, adjacent to the abutments, and along the banks adjacent to the bridges. Outside of the banks, developed land includes areas that are paved, graded, hardpack dirt, and gravel access routes. These areas were generally devoid of substantial vegetation cover but contained small patches of non-native vegetation. Species observed includes were the same as observed in the Wild Oats and Annual Brome Grassland (Photo 6, Appendix D).

3.8 Areas Not Meeting the Regulatory Definition of Section 404/401 Wetlands and Waters

In general, areas that were not considered to be wetlands were not dominated by hydrophytic vegetation and did not exhibit hydrology indicators. Approximately 10.39 acres of the study area met none of the regulatory definitions of jurisdictional waters or jurisdictional habitats including the following habitat/land cover types: annual grassland (3.92 acres), coast live oak woodland (0.38 acres), ornamental woodland (1.25 acres) and developed land (4.84 acres) (Figure 6).

Ornamental Woodland. Ornamental woodland includes lands that have been planted with landscaping and are maintained on an ongoing basis. Such landscaping may include native and non-native plantings. Within the study area, ornamental woodland is found along both sides of the tracks in the eastern reach of the study area and includes the City of San Jose's Fuller Avenue Park. Trees and shrubs observed included black locust (*Robinia pseudoacacia*), Chinese pistache (*Pistacia chinensis*), Australian pine (*Casuarina equisetifolia*), Canary Island pine (*Pinus canariensis*), and cotoneaster (*Cotoneaster franchetii*). Ornamental woodland is found along the top of bank of the flood control basin and McClellan Avenue in a small landscaped neighborhood park. Trees and shrubs observed included blue elderberry (*Sambucus nigra* ssp. *caerulea*), valley oak (*Quercus lobata*), and California coffeeberry (*Frangula californica*).

Coast Live Oak Woodland. Woodland habitat dominated by coast live oak (*Quercus agrifolia*) occurs in two areas within the study area. A small amount of the oak woodland canopy overhangs the top of bank in the northwest corner of the study area, but the trees are rooted outside of the top of bank; therefore, they were mapped as part of the Coast Live Oak Woodland. Plants observed in the understory were the same as those observed in the Wild Oats and Annual Brome Grassland.



Figure 6 Vegetation Communities Map

Feet 400

Guadalupe River Bridge Replacement Project

Study Area (17.06 acres)
Freshwater Perennial Marsh (1.61 acres)
Seasonal Wetland (1.33 acres)
Aquatic Habitat (0.73 acres)

Fremont Cottonwood Forest (2.13 acres)

MIG



Section 404/401 Waters (Up to OHWM) **1.45 acres**

Section 404/401 Wetlands 2.94 acres

Section 401 (up to top of bank) 1.66 acres

CDFW Jurisdictional Habitats 6.67 acres

- Study Area (17.06 acres)
- CII Ordinary High Water Mark (OHWM)
- Top of Bank (TOB)
- **OHWM Sample Point** •
- Wetland Sample Point •
- Photo Point
- Culvert Ο



- Freshwater Perennial Marsh (1.61 acres)
- Seasonal Wetland (1.33 acres)
- Open Water River Channel (0.73 acres)
- Fremont Cottonwood Forest (2.13 acres)
- Coast Live Oak Woodland (0.38 acres)
 - Wild Oats and Annual Brome Grassland (4.59 acres)
- Ornamental Woodland (1.25 acres)
- Developed Land (5.04 acres)



Figure 7 Preliminary Identification of Waters of the U.S./State

Guadalupe River Bridge Replacement Project

MT

MT2



4. References

- Baldwin, Goldman, Keil, Patterson, Ronatti, and Wilkin (eds.) 2012. The Jepson Manual: Vascular Plants of California. 2nd Edition. University of California Press, Berkeley, California.
- [CDFW] California Department of Fish and Wildlife. 2020. VegCAMP Natural Communities Lists. Accessed March 2020 from https://www.wildlife.ca.gov/data/vegcamp/natural-communities.
- Cowardin, L. M., V. Carter, F. C. Golet, E. T. LaRoe. 1979. Classification of Wetlands and Deepwater Habitats of the United States. U. S. Department of the Interior, Fish and Wildlife Service, Washington, D.C. Jamestown, ND: Northern Prairie Wildlife Research Center Home Page. (Version 04DEC98).
- Environmental Laboratory. 1987. U.S. Corps of Engineers Wetlands Delineation Manual. Department of the Army.
- [FEMA] Federal Emergency Management Agency. 2019. Flood Insurance Rate Map 06085C0234H, 05/18/2009.
- Google Inc. 2019, 2020. Google Earth Pro (Version 7.1.5.1557) [Software]. Available from earth.google.com.
- Lichvar, R. W., D. L. Banks, W. N. Kirchner, and N. C. Melvin. 2016. Western Mountains, Valleys, and Coast 2016 Regional Plant List. The National Wetland Plant List: 2016 wetland ratings. Phytoneuron 2016-30: 1-17. Prepared for the U.S. Army Corps of Engineers.
- Munsell. 2009. Soil Color Charts, Munsell Color X-rite. Grand Rapids, Michigan.
- [NRCS] Natural Resources Conservation Service. 2010. Field Indicators of Hydric Soils in the U.S.: A Guide for Identifying and Delineating Hydric Soils (Version 7.0). U.S. Department of Agriculture. Prepared with the National Technical Committee for Hydric Soils.
- [NRCS] Natural Resources Conservation Service. 2019a. Web Soil Survey. U.S. Department of Agriculture. Accessed November 2019 from http://websoilsurvey.nrcs.usda.gov.
- [NRCS] National Resource Conservation Service. 2019b. National Hydric Soils List. Accessed November 2019 from http://www.nrcs.usda.gov/wps/portal/nrcs/main/soils/use/hydric/.
- [NWI] National Wetlands Inventory. 2019. Wetlands Mapper. U.S. Fish and Wildlife Service Accessed November 2019 from http://www.fws.gov/wetlands/Wetlands-Mapper.html.
- PRISM Climate Group. 2019. Online PRISM Data Explorer. Oregon State University, Corvallis, Oregon. Accessed November 2019 from http://www.prism.oregonstate.edu/.
- [UCSB] University of California Santa Barbara Library. 2020. Digital Aerial Photography Collections. Accessed November 2020 from https://www.library.ucsb.edu/src/airphotos
- [USACE] U.S. Army Corps of Engineers. 2004. Review of Ordinary High Water Mark Indicators for Delineating Arid Streams in the Southwestern United States. January 2004.

- [USACE] U.S. Army Corps of Engineers. 2007. Review and Synopsis of Natural and Human Controls on Fluvial Channel Processes in the Arid West. September 2007.
- [USACE] U.S. Army Corps of Engineers. 2008a. Regional Supplement to the Corps of Engineers Wetland Delineation Manual: Western Mountains, Valleys, and Coast Region. May 2010. U.S. Army Engineer Research and Development Center.
- [USACE] U.S. Army Corps of Engineers. 2008b. A Field Guide to the Identification of the Ordinary High-Water Mark (OHWM) in the Arid West Region of the Western United States. August 2008. U.S. Army Engineer Research and Development Center.
- [USACE] U.S. Army Corps of Engineers. 2016a. Updated Map and Drawing Standards for the South Pacific Division Regulatory Program.
- [USACE] U.S. Army Corps of Engineers. 2016b. Information Needed for Verification of Corps Jurisdiction, San Francisco District. Revised April 2016. https://www.spn.usace.army.mil/Portals/68/docs/regulatory/2%20-%20Info%20Req.pdf
- [USGS] U.S. Geological Survey. 2019. Water Resources of the United States. Accessed November 2019 from https://water.usgs.gov/GIS/huc_name.html#Region18

Appendix A: Soil Report for the Study Area



United States Department of Agriculture

Natural Resources Conservation

Service

A product of the National Cooperative Soil Survey, a joint effort of the United States Department of Agriculture and other Federal agencies, State agencies including the Agricultural Experiment Stations, and local participants

Custom Soil Resource Report for Santa Clara Area, California, Western Part

Guadalupe River Bridge Replacement Project



Preface

Soil surveys contain information that affects land use planning in survey areas. They highlight soil limitations that affect various land uses and provide information about the properties of the soils in the survey areas. Soil surveys are designed for many different users, including farmers, ranchers, foresters, agronomists, urban planners, community officials, engineers, developers, builders, and home buyers. Also, conservationists, teachers, students, and specialists in recreation, waste disposal, and pollution control can use the surveys to help them understand, protect, or enhance the environment.

Various land use regulations of Federal, State, and local governments may impose special restrictions on land use or land treatment. Soil surveys identify soil properties that are used in making various land use or land treatment decisions. The information is intended to help the land users identify and reduce the effects of soil limitations on various land uses. The landowner or user is responsible for identifying and complying with existing laws and regulations.

Although soil survey information can be used for general farm, local, and wider area planning, onsite investigation is needed to supplement this information in some cases. Examples include soil quality assessments (http://www.nrcs.usda.gov/wps/portal/nrcs/main/soils/health/) and certain conservation and engineering applications. For more detailed information, contact your local USDA Service Center (https://offices.sc.egov.usda.gov/locator/app?agency=nrcs) or your NRCS State Soil Scientist (http://www.nrcs.usda.gov/wps/portal/nrcs/detail/soils/contactus/? cid=nrcs142p2_053951).

Great differences in soil properties can occur within short distances. Some soils are seasonally wet or subject to flooding. Some are too unstable to be used as a foundation for buildings or roads. Clayey or wet soils are poorly suited to use as septic tank absorption fields. A high water table makes a soil poorly suited to basements or underground installations.

The National Cooperative Soil Survey is a joint effort of the United States Department of Agriculture and other Federal agencies, State agencies including the Agricultural Experiment Stations, and local agencies. The Natural Resources Conservation Service (NRCS) has leadership for the Federal part of the National Cooperative Soil Survey.

Information about soils is updated periodically. Updated information is available through the NRCS Web Soil Survey, the site for official soil survey information.

The U.S. Department of Agriculture (USDA) prohibits discrimination in all its programs and activities on the basis of race, color, national origin, age, disability, and where applicable, sex, marital status, familial status, parental status, religion, sexual orientation, genetic information, political beliefs, reprisal, or because all or a part of an individual's income is derived from any public assistance program. (Not all prohibited bases apply to all programs.) Persons with disabilities who require

alternative means for communication of program information (Braille, large print, audiotape, etc.) should contact USDA's TARGET Center at (202) 720-2600 (voice and TDD). To file a complaint of discrimination, write to USDA, Director, Office of Civil Rights, 1400 Independence Avenue, S.W., Washington, D.C. 20250-9410 or call (800) 795-3272 (voice) or (202) 720-6382 (TDD). USDA is an equal opportunity provider and employer.
Contents

Preface	2
How Soil Surveys Are Made	5
Soil Map	8
Soil Map	9
Legend	10
Map Unit Legend	11
Map Unit Descriptions	11
Santa Clara Area, California, Western Part	13
165-Urbanland-Campbell complex, 0 to 2 percent slopes, protected	13
References	15

How Soil Surveys Are Made

Soil surveys are made to provide information about the soils and miscellaneous areas in a specific area. They include a description of the soils and miscellaneous areas and their location on the landscape and tables that show soil properties and limitations affecting various uses. Soil scientists observed the steepness, length, and shape of the slopes; the general pattern of drainage; the kinds of crops and native plants; and the kinds of bedrock. They observed and described many soil profiles. A soil profile is the sequence of natural layers, or horizons, in a soil. The profile extends from the surface down into the unconsolidated material in which the soil formed or from the surface down to bedrock. The unconsolidated material is devoid of roots and other living organisms and has not been changed by other biological activity.

Currently, soils are mapped according to the boundaries of major land resource areas (MLRAs). MLRAs are geographically associated land resource units that share common characteristics related to physiography, geology, climate, water resources, soils, biological resources, and land uses (USDA, 2006). Soil survey areas typically consist of parts of one or more MLRA.

The soils and miscellaneous areas in a survey area occur in an orderly pattern that is related to the geology, landforms, relief, climate, and natural vegetation of the area. Each kind of soil and miscellaneous area is associated with a particular kind of landform or with a segment of the landform. By observing the soils and miscellaneous areas in the survey area and relating their position to specific segments of the landform, a soil scientist develops a concept, or model, of how they were formed. Thus, during mapping, this model enables the soil scientist to predict with a considerable degree of accuracy the kind of soil or miscellaneous area at a specific location on the landscape.

Commonly, individual soils on the landscape merge into one another as their characteristics gradually change. To construct an accurate soil map, however, soil scientists must determine the boundaries between the soils. They can observe only a limited number of soil profiles. Nevertheless, these observations, supplemented by an understanding of the soil-vegetation-landscape relationship, are sufficient to verify predictions of the kinds of soil in an area and to determine the boundaries.

Soil scientists recorded the characteristics of the soil profiles that they studied. They noted soil color, texture, size and shape of soil aggregates, kind and amount of rock fragments, distribution of plant roots, reaction, and other features that enable them to identify soils. After describing the soils in the survey area and determining their properties, the soil scientists assigned the soils to taxonomic classes (units). Taxonomic classes are concepts. Each taxonomic class has a set of soil characteristics with precisely defined limits. The classes are used as a basis for comparison to classify soils systematically. Soil taxonomy, the system of taxonomic classification used in the United States, is based mainly on the kind and character of soil properties and the arrangement of horizons within the profile. After the soil

scientists classified and named the soils in the survey area, they compared the individual soils with similar soils in the same taxonomic class in other areas so that they could confirm data and assemble additional data based on experience and research.

The objective of soil mapping is not to delineate pure map unit components; the objective is to separate the landscape into landforms or landform segments that have similar use and management requirements. Each map unit is defined by a unique combination of soil components and/or miscellaneous areas in predictable proportions. Some components may be highly contrasting to the other components of the map unit. The presence of minor components in a map unit in no way diminishes the usefulness or accuracy of the data. The delineation of such landforms and landform segments on the map provides sufficient information for the development of resource plans. If intensive use of small areas is planned, onsite investigation is needed to define and locate the soils and miscellaneous areas.

Soil scientists make many field observations in the process of producing a soil map. The frequency of observation is dependent upon several factors, including scale of mapping, intensity of mapping, design of map units, complexity of the landscape, and experience of the soil scientist. Observations are made to test and refine the soil-landscape model and predictions and to verify the classification of the soils at specific locations. Once the soil-landscape model is refined, a significantly smaller number of measurements of individual soil properties are made and recorded. These measurements may include field measurements, such as those for color, depth to bedrock, and texture, and laboratory measurements, such as those for content of sand, silt, clay, salt, and other components. Properties of each soil typically vary from one point to another across the landscape.

Observations for map unit components are aggregated to develop ranges of characteristics for the components. The aggregated values are presented. Direct measurements do not exist for every property presented for every map unit component. Values for some properties are estimated from combinations of other properties.

While a soil survey is in progress, samples of some of the soils in the area generally are collected for laboratory analyses and for engineering tests. Soil scientists interpret the data from these analyses and tests as well as the field-observed characteristics and the soil properties to determine the expected behavior of the soils under different uses. Interpretations for all of the soils are field tested through observation of the soils in different uses and under different levels of management. Some interpretations are modified to fit local conditions, and some new interpretations are developed to meet local needs. Data are assembled from other sources, such as research information, production records, and field experience of specialists. For example, data on crop yields under defined levels of management are assembled from farm records and from field or plot experiments on the same kinds of soil.

Predictions about soil behavior are based not only on soil properties but also on such variables as climate and biological activity. Soil conditions are predictable over long periods of time, but they are not predictable from year to year. For example, soil scientists can predict with a fairly high degree of accuracy that a given soil will have a high water table within certain depths in most years, but they cannot predict that a high water table will always be at a specific level in the soil on a specific date.

After soil scientists located and identified the significant natural bodies of soil in the survey area, they drew the boundaries of these bodies on aerial photographs and

identified each as a specific map unit. Aerial photographs show trees, buildings, fields, roads, and rivers, all of which help in locating boundaries accurately.

Soil Map

The soil map section includes the soil map for the defined area of interest, a list of soil map units on the map and extent of each map unit, and cartographic symbols displayed on the map. Also presented are various metadata about data used to produce the map, and a description of each soil map unit.



	MAP LI	EGEND		MAP INFORMATION
Area of Interest (AOI)		000	Spoil Area	The soil surveys that comprise your AOI were mapped at
Are	ea of Interest (AOI)	۵	Stony Spot	1:24,000.
Soils	il Man Unit Dalvaana	Ø	Very Stony Spot	Warning: Soil Map may not be valid at this scale.
50	il Map Unit Polygons	Ŷ	Wet Spot	
~ 50		Δ	Other	Enlargement of maps beyond the scale of mapping can cause
	Soil Map Unit Points		Special Line Features	line placement. The maps do not show the small areas of
Special Poin	t Features	Water Feat	tures	contrasting soils that could have been shown at a more detailed scale
Ba Ba	rrow Pit	\sim	Streams and Canals	
	now r it	Transporta	ation	Please rely on the bar scale on each map sheet for map
		+++ R	Rails	measurements.
	bsed Depression	~	Interstate Highways	Source of Map: Natural Resources Conservation Service
K Gra	Gravelly Spot	✓ US Routes✓ Major Roads	US Routes	Web Soil Survey URL:
• Gra	avelly Spot		Coordinate System: Web Mercator (EPSG:3857)	
🙆 Lai	ndfill	~	Local Roads	Maps from the Web Soil Survey are based on the Web Mercator
A Lav	va Flow	Backgrour		projection, which preserves direction and shape but distorts distance and area. A projection that preserves area, such as the
🚲 Ma	arsh or swamp	No.	Aerial Photography	Albers equal-area conic projection, should be used if more
🙊 Mir	ne or Quarry			accurate calculations of distance or area are required.
Mis	scellaneous Water		This product is generated from the USDA-NRCS certified data as	
O Pe	rennial Water			of the version date(s) listed below.
👽 Ro	ck Outcrop			Soil Survey Area: Santa Clara Area, California, Western Part
🕂 Sa	line Spot			Survey Area Data: Version 9, May 29, 2020
sa Sa	ndy Spot			Soil map units are labeled (as space allows) for map scales
e Se	verely Eroded Spot			1:50,000 or larger.
👌 Sir	hhole			Date(s) aerial images were photographed: Apr 13, 2010_Apr
🚡 Slie	de or Slip			23, 2019
ø So	dic Spot			The orthophoto or other base map on which the soil lines were compiled and digitized probably differs from the background imagery displayed on these maps. As a result, some minor shifting of map unit boundaries may be evident

Map Unit Legend

Map Unit Symbol	Map Unit Name	Acres in AOI	Percent of AOI
165	Urbanland-Campbell complex, 0 to 2 percent slopes, protected	17.9	100.0%
Totals for Area of Interest		17.9	100.0%

Map Unit Descriptions

The map units delineated on the detailed soil maps in a soil survey represent the soils or miscellaneous areas in the survey area. The map unit descriptions, along with the maps, can be used to determine the composition and properties of a unit.

A map unit delineation on a soil map represents an area dominated by one or more major kinds of soil or miscellaneous areas. A map unit is identified and named according to the taxonomic classification of the dominant soils. Within a taxonomic class there are precisely defined limits for the properties of the soils. On the landscape, however, the soils are natural phenomena, and they have the characteristic variability of all natural phenomena. Thus, the range of some observed properties may extend beyond the limits defined for a taxonomic class. Areas of soils of a single taxonomic class rarely, if ever, can be mapped without including areas of other taxonomic classes. Consequently, every map unit is made up of the soils or miscellaneous areas for which it is named and some minor components that belong to taxonomic classes other than those of the major soils.

Most minor soils have properties similar to those of the dominant soil or soils in the map unit, and thus they do not affect use and management. These are called noncontrasting, or similar, components. They may or may not be mentioned in a particular map unit description. Other minor components, however, have properties and behavioral characteristics divergent enough to affect use or to require different management. These are called contrasting, or dissimilar, components. They generally are in small areas and could not be mapped separately because of the scale used. Some small areas of strongly contrasting soils or miscellaneous areas are identified by a special symbol on the maps. If included in the database for a given area, the contrasting minor components are identified in the map unit descriptions along with some characteristics of each. A few areas of minor components may not have been observed, and consequently they are not mentioned in the descriptions, especially where the pattern was so complex that it was impractical to make enough observations to identify all the soils and miscellaneous areas on the landscape.

The presence of minor components in a map unit in no way diminishes the usefulness or accuracy of the data. The objective of mapping is not to delineate pure taxonomic classes but rather to separate the landscape into landforms or landform segments that have similar use and management requirements. The delineation of such segments on the map provides sufficient information for the development of resource plans. If intensive use of small areas is planned, however,

onsite investigation is needed to define and locate the soils and miscellaneous areas.

An identifying symbol precedes the map unit name in the map unit descriptions. Each description includes general facts about the unit and gives important soil properties and qualities.

Soils that have profiles that are almost alike make up a *soil series*. Except for differences in texture of the surface layer, all the soils of a series have major horizons that are similar in composition, thickness, and arrangement.

Soils of one series can differ in texture of the surface layer, slope, stoniness, salinity, degree of erosion, and other characteristics that affect their use. On the basis of such differences, a soil series is divided into *soil phases*. Most of the areas shown on the detailed soil maps are phases of soil series. The name of a soil phase commonly indicates a feature that affects use or management. For example, Alpha silt loam, 0 to 2 percent slopes, is a phase of the Alpha series.

Some map units are made up of two or more major soils or miscellaneous areas. These map units are complexes, associations, or undifferentiated groups.

A *complex* consists of two or more soils or miscellaneous areas in such an intricate pattern or in such small areas that they cannot be shown separately on the maps. The pattern and proportion of the soils or miscellaneous areas are somewhat similar in all areas. Alpha-Beta complex, 0 to 6 percent slopes, is an example.

An *association* is made up of two or more geographically associated soils or miscellaneous areas that are shown as one unit on the maps. Because of present or anticipated uses of the map units in the survey area, it was not considered practical or necessary to map the soils or miscellaneous areas separately. The pattern and relative proportion of the soils or miscellaneous areas are somewhat similar. Alpha-Beta association, 0 to 2 percent slopes, is an example.

An *undifferentiated group* is made up of two or more soils or miscellaneous areas that could be mapped individually but are mapped as one unit because similar interpretations can be made for use and management. The pattern and proportion of the soils or miscellaneous areas in a mapped area are not uniform. An area can be made up of only one of the major soils or miscellaneous areas, or it can be made up of all of them. Alpha and Beta soils, 0 to 2 percent slopes, is an example.

Some surveys include *miscellaneous areas*. Such areas have little or no soil material and support little or no vegetation. Rock outcrop is an example.

Santa Clara Area, California, Western Part

165—Urbanland-Campbell complex, 0 to 2 percent slopes, protected

Map Unit Setting

National map unit symbol: 1qsvl Elevation: 0 to 240 feet Mean annual precipitation: 14 to 24 inches Mean annual air temperature: 57 to 61 degrees F Frost-free period: 275 to 325 days Farmland classification: Not prime farmland

Map Unit Composition

Urban land: 70 percent Campbell, protected, and similar soils: 20 percent Minor components: 10 percent Estimates are based on observations, descriptions, and transects of the mapunit.

Description of Urban Land

Setting

Landform: Alluvial fans Landform position (three-dimensional): Talf Down-slope shape: Linear Across-slope shape: Linear Parent material: Disturbed and human-transported material

Description of Campbell, Protected

Setting

Landform: Alluvial fans Landform position (three-dimensional): Talf Down-slope shape: Linear Across-slope shape: Linear Parent material: Alluvium derived from metamorphic and sedimentary rock and/or alluvium derived from metavolcanics

Typical profile

Ap - 0 to 10 inches: silt loam A1 - 10 to 24 inches: silt loam A2 - 24 to 31 inches: silty clay loam A3 - 31 to 38 inches: silty clay loam 2A - 38 to 51 inches: silty clay loam 2Bw1 - 51 to 71 inches: silty clay 2Bw2 - 71 to 79 inches: silty clay

Properties and qualities

Slope: 0 to 2 percent
Depth to restrictive feature: More than 80 inches
Drainage class: Moderately well drained
Runoff class: Very low
Capacity of the most limiting layer to transmit water (Ksat): Moderately low to moderately high (0.06 to 0.20 in/hr)
Depth to water table: More than 80 inches
Frequency of flooding: None

Frequency of ponding: None *Calcium carbonate, maximum content:* 15 percent *Maximum salinity:* Nonsaline to slightly saline (1.0 to 4.0 mmhos/cm) *Sodium adsorption ratio, maximum:* 5.0 *Available water capacity:* High (about 10.4 inches)

Interpretive groups

Land capability classification (irrigated): 1 Land capability classification (nonirrigated): 3s Hydrologic Soil Group: C Hydric soil rating: No

Minor Components

Clear lake

Percent of map unit: 5 percent Landform: Basin floors Landform position (three-dimensional): Talf Down-slope shape: Linear Across-slope shape: Linear Hydric soil rating: Yes

Newpark

Percent of map unit: 5 percent Landform: Alluvial fans Landform position (three-dimensional): Talf Down-slope shape: Linear Across-slope shape: Linear Hydric soil rating: No

References

American Association of State Highway and Transportation Officials (AASHTO). 2004. Standard specifications for transportation materials and methods of sampling and testing. 24th edition.

American Society for Testing and Materials (ASTM). 2005. Standard classification of soils for engineering purposes. ASTM Standard D2487-00.

Cowardin, L.M., V. Carter, F.C. Golet, and E.T. LaRoe. 1979. Classification of wetlands and deep-water habitats of the United States. U.S. Fish and Wildlife Service FWS/OBS-79/31.

Federal Register. July 13, 1994. Changes in hydric soils of the United States.

Federal Register. September 18, 2002. Hydric soils of the United States.

Hurt, G.W., and L.M. Vasilas, editors. Version 6.0, 2006. Field indicators of hydric soils in the United States.

National Research Council. 1995. Wetlands: Characteristics and boundaries.

Soil Survey Division Staff. 1993. Soil survey manual. Soil Conservation Service. U.S. Department of Agriculture Handbook 18. http://www.nrcs.usda.gov/wps/portal/ nrcs/detail/national/soils/?cid=nrcs142p2_054262

Soil Survey Staff. 1999. Soil taxonomy: A basic system of soil classification for making and interpreting soil surveys. 2nd edition. Natural Resources Conservation Service, U.S. Department of Agriculture Handbook 436. http://www.nrcs.usda.gov/wps/portal/nrcs/detail/national/soils/?cid=nrcs142p2_053577

Soil Survey Staff. 2010. Keys to soil taxonomy. 11th edition. U.S. Department of Agriculture, Natural Resources Conservation Service. http://www.nrcs.usda.gov/wps/portal/nrcs/detail/national/soils/?cid=nrcs142p2_053580

Tiner, R.W., Jr. 1985. Wetlands of Delaware. U.S. Fish and Wildlife Service and Delaware Department of Natural Resources and Environmental Control, Wetlands Section.

United States Army Corps of Engineers, Environmental Laboratory. 1987. Corps of Engineers wetlands delineation manual. Waterways Experiment Station Technical Report Y-87-1.

United States Department of Agriculture, Natural Resources Conservation Service. National forestry manual. http://www.nrcs.usda.gov/wps/portal/nrcs/detail/soils/ home/?cid=nrcs142p2 053374

United States Department of Agriculture, Natural Resources Conservation Service. National range and pasture handbook. http://www.nrcs.usda.gov/wps/portal/nrcs/ detail/national/landuse/rangepasture/?cid=stelprdb1043084

United States Department of Agriculture, Natural Resources Conservation Service. National soil survey handbook, title 430-VI. http://www.nrcs.usda.gov/wps/portal/ nrcs/detail/soils/scientists/?cid=nrcs142p2_054242

United States Department of Agriculture, Natural Resources Conservation Service. 2006. Land resource regions and major land resource areas of the United States, the Caribbean, and the Pacific Basin. U.S. Department of Agriculture Handbook 296. http://www.nrcs.usda.gov/wps/portal/nrcs/detail/national/soils/? cid=nrcs142p2_053624

United States Department of Agriculture, Soil Conservation Service. 1961. Land capability classification. U.S. Department of Agriculture Handbook 210. http://www.nrcs.usda.gov/Internet/FSE_DOCUMENTS/nrcs142p2_052290.pdf

Appendix B: Plants Observed in the Study Area

Common Name	Scientific Name	Wetland Indicator Status ¹
American wild carrot	Daucus pusillus	NI
Arroyo willow	Salix Iasiolepis	FACW
Australian pine	Casuarina equisetifolia	FACU
Boxelder	Acer negundo	FACW
Bird's foot trefoil	Lotus corniculatus	FAC
Black locust	Robinia pseudoacacia	FACU
Black mustard	Brassica nigra	NI
Blue elderberry	Sambucus nigra ssp. caerulea	FACU
Bristly ox-tongue	Helminthotheca echioides	FAC
Broadleaf cattail	Typha latifolia	OBL
California coffeeberry	Frangula californica	NI
California mugwort	Artemisia douglasiana	FAC
Canada horseweed	Erigeron canadensis	FACU
Canary Island pine	Pinus canariensis	NI
Chinese pistache	Pistacia chinensis	NI
Coast live oak	Quercus agrifolia	NI
Common mallow	Malva neglecta	NI
Cotoneaster	Cotoneaster franchetii	NI
Curly dock	Rumix crispis	FAC
Dallis grass	Paspalum dilatatum	FAC
Devil's beggartick	Bidens frondosa	FACW
Dotted smartweed	Persicaria punctata	OBL
English ivy	Hedera helix	FACU
English plantain	Plantago lanceolata	FAC
Field bindweed	Convolvulus arvensis	NI
Fremont cottonwood	Populus fremontii	NI (considered riparian)
Fringed willowherb	Epilobium ciliatum	FACW
Giant reed	Arundo donax	FACW
Harding grass	Phalaris aquatica	FACU
Hardstem bulrush	Schoenoplectus acutus	OBL
Heliotrope	Heliotropium curassavicum	FACU
Himalayan blackberry	Rubus armeniacus	FAC
Italian rye grass	Festuca perennis	FAC
Italian thistle	Carduus pycnocephalus	NI
Mediterranean barley	Hordeum murinum	FACU

Mule fat	Baccharis salicifolia	FAC
Poison hemlock	Conium maculatum	FACW
Prickly lettuce	Lactuca serriola	FACU
Rabbitsfoot grass	Polypogon monspeliensis	FACW
Red willow	Salix laevigata	FACW
Rough cocklebur	Xanthium strumarium	FAC
Ripgut brome	Bromus diandrus	NI
Russian thistle	Salsola tragus	FACU
Seaside barley	Hordeum marinum	FAC
Spearmint	Mentha spicata	FACW
Sweet fennel	Foeniculum vulgare	NI
Tall flatsedge	Cyperus eragrostis	FACW
Prostrate knotweed	Polygonum aviculare	FAC
Puncture vine	Tribulus terrestris	NI
Smilo grass	Stipa miliacea	NI
Valley oak	Quercus lobata	FACU
Water primrose	Ludwigia peploides	OBL
Wild oat	Avena fatua	NI
White horehound	Marrubium vulgare	FACU
White sweetclover	Melilotus albus	NI
Wild radish	Raphanus sativus	NI

Notes:

¹Wetland Indicator Status obtained from Lichvar et al. (2016)

Wetland Indicator Status Key:

OBL = Obligate wetland species, occur almost always in wetlands (>99% probability).

FACW = Facultative Wetland species, usually occur in wetlands (67 to 99% probability), but occasionally found in non-wetlands.

FAC = Facultative species, equally likely to occur in wetlands or non-wetlands (34 to 66% probability).

FACU = Facultative Upland, usually occur in non-wetlands (67% to 99%), but occasionally found in wetlands.

UPL = Obligate Upland species, occur almost always in non-wetlands (>99% probability).

NI = Non-Indicator, not present on list. Considered to be an upland species unless otherwise indicated.

Appendix C: USACE Western Mountains, Valley and Coast Wetland Data Forms

WETLAND DETERMINATION DATA FORM - Arid West Region

Project/Site: Guadalupe River Bridge Replacement Project	City/County: San Jose/San	Sampling Date: 14-Aug-20		g-20	
Applicant/Owner: Peninsula Corridor Joint Powers Board	Sta	te: CA	Sampling Poin	it:	SP1
Investigator(s): DWG, AB	Section, Township, Ra	nge: S	т	R	
Landform (hillslope, terrace, etc.): Floodplain	Local relief (concave, co	onvex, none): flat		Slope:	% / °
Subregion (LRR): LRR C Lat.:	37.317311 Long.: -121.887825			Datum: WGS84	
Soil Map Unit Name: <u>165 – Urbanland-Campbell complex</u> , 0 to 2 perce	nt slopes	NWI o	classification: P	'EM1E	
Are climatic/hydrologic conditions on the site typical for this time of ye	ear? Yes $ullet$ No $igodot$	(If no, expla	in in Remarks.)	
Are Vegetation 🗹 , Soil 🗌 , or Hydrology 🗌 significant	tly disturbed? Are "No	rmal Circumstanc	es" present?	Yes \bigcirc	No 🖲
Are Vegetation 🗌 , Soil 🗌 , or Hydrology 🗌 naturally	problematic? (If need	led, explain any a	nswers in Rem	arks.)	

Summary of Findings - Attach site map showing sampling point locations, transects, important features, etc.

Hydrophytic Vegetation Present?	Yes 🖲	No O	Is the Sampled Area		
Hydric Soil Present?	Yes 🖲	No 🔿		Yes 🔍 No 🔿	
Wetland Hydrology Present?	Yes 🖲	No 🔿	within a wetland?		
Remarks: Area recently mowed.					

Dominant

VEGETATION - Use scientific names of plants.

		-Species?		
Tree Stratum (Plot size:	Absolute	Rel.Strat.	Indicator	Dominance Test worksheet:
			Status	Number of Dominant Species
l,				That are OBL, FACW, or FAC: (A)
2	0	0.0%		Total Number of Dominant
3	0	0.0%		Species Across All Strata: 1 (B)
4.	0	0.0%		
		- Total Cove		Percent of dominant Species
Sapling/Shrub Stratum (Plot size:)			-	That Are OBL, FACW, or FAC:
1	0	0.0%		
1				Prevalence Index worksheet:
2	0	0.0%		Total % Cover of: Multiply by:
3	0	0.0%		OBL species $0 \times 1 = 0$
4.	0	0.0%		FACW species $0 \times 2 = 0$
5.	0	0.0%		EAC species $30 \times 3 = 90$
	0	= Total Cove	er	FACU species $0 \times 4 = 0$
<u>Herb Stratum</u> (Plot size: <u>5 x 5</u>)				
1. Helminthotheca echioides	30	✔ 96.8%	FAC	$\frac{1}{1}$
2. Convolvulus arvensis	1	3.2%	UPL	Column Totals: <u>31</u> (A) <u>95</u> (B)
3.	0	0.0%		Prevalence Index = $B/A = 3.065$
4.	0	0.0%		Hydrophytic Vegetation Indicators
5.	0	0.0%		\mathbf{V} Dominance Test is > 50%
6.	0	0.0%		
7.				
8.		0.0%		Morphological Adaptations ¹ (Provide supporting
9				Gata in Remarks of on a separate sneet)
10.		0.0%		Problematic Hydrophytic Vegetation ' (Explain)
11.				
				¹ Indicators of hydric soil and wetland hydrology must
	31	= Total Cove	er	be present, unless disturbed or problematic.
Woodv Vine Stratum (Plot size:)				
1	0	0.0%		
2.	0	0.0%		Hydrophytic
	0	= Total Cove	er	Vegetation Present? Yes • No ·
% Bare Ground in Herb Stratum: 0 % Co	ver of Biotio	c Crust ₍₎		
Remarks:		<u> </u>		1

No bare ground, remaining area covered in a thick layer of thatch (~1 inch). Vigor response by bristly ox-tongue.

*Indicator suffix = National status or professional decision assigned because Regional status not defined by FWS.

Soil

Sampling Point: SP1

V VII								Sampi		
Profile Descri	iption: (Describ	e to the depth	needed t	o document	the indi	cator or co	nfirm the a	absence of indicator	rs.)	
Depth (inches)	Mat	rix	Cala	Red (moist)	lox Featu	ires Turna ¹		Touturo	Domostra	
		/2 95	 5VD	4/6	<u>-70</u> 5		_ <u>LOC-</u>	Silty Clay Loam	Soil moist	
0-10										
			-							
				-	-	-				
					-					
	. <u> </u>									
¹ Type: C=Cond	centration. D=Dep	oletion. RM=Red	uced Matri	x, CS=Covere	d or Coate	ed Sand Grai	ins ² Locat	ion: PL=Pore Lining.	1=Matrix	
Hydric Soil I	ndicators: (App	licable to all L	RRs, unle	ss otherwis	e noted.)		Indicators for P	roblematic Hydric Soils. ³	
	A1)			andy Redox ((S5)			1 cm Muck (A	9) (LRR C)	
Black Histi	(A2)			tripped Matri	x (S6)			2 cm Muck (A	10) (LRR B)	
Hvdrogen	Sulfide (A4)			oamy Mucky	Mineral (F	1)		Reduced Verti	c (F18)	
Stratified L	_avers (A5) (LRR (C)		oamy Gleyed	Matrix (F2	2)		Red Parent Ma	aterial (TF2)	
1 cm Muck	(A9) (LRR D)	- /		Pepleted Matr	ix (F3)			Other (Explain	n in Remarks)	
Depleted E	Below Dark Surfac	e (A11)		edox Dark Su	urface (F6))				
Thick Dark	Surface (A12)			epieted Dark	Surface (F7)				
Sandy Muc	ck Mineral (S1)		▼ F	edox depress				³ Indicators of hyd	drophytic vegetation and	
Sandy Gley	yed Matrix (S4)				F9)			wetland hydrol	ogy must be present.	
Restrictive La	yer (if present)	:								
Type:										
Depth (inch	nes):							Hydric Soil Preser	it? Yes 🖲 No 🔾	
Remarks:										
Hydrology	1									
Wetland Hydr	rology Indicator	·c'								
		s.	od, chool	all that an	-			Cocondon	Indicators (2 or more requ	irod)
	ator (A1)	n of one requir		Salt Cruct (F	<u>))))</u> 211)				Indicators (2 or more required)	irea)
High Wate	ater (AI) r Table (A2)			Biotic Crust	(B12)				diks (D1) (Riverine)	
	(A3)				ortebrates	(B13)			nceits (B2) (Riverine)	
Water Mar	(Nonriveri	ne)		Hydrogen Si	ulfide Odo	(D13) r (C1)			Patterns (B10)	
Sediment I	Denosits (B2) (No	nriverine)			izosnheres	along Livin	a Roots (C3) Dru Sea	con Water Table (C2)	
	sits (B3) (Nonerive	erine)		Presence of	Reduced I	ron (C4)	g 10003 (05	Cravfish	Burrows (C8)	
Surface So	oil Cracks (B6)	child)		Recent Iron	Reduction	in Plowed 9	Soils (C6)	Saturati	on Visible on Aerial Imagery (C	9)
✓ Inundation	n Visible on Aerial	Imagery (B7)		Thin Muck S	urface (C7	')	()		Aquitard (D3)	
Water-Stai	ined Leaves (B9)			Other (Expla	ain in Rem	, arks)		FAC-neu	tral Test (D5)	
						/				
Field Observa	Procont?	Yes 🔿 No	\odot	Denth (inc	hes).					
	Present?			Depth (inc	iics).		_			
Water Table Pro	esent?	Yes \bigcirc No	lacksquare	Depth (inc	hes):		Wetla	nd Hydrology Prese	nt? Yes 🖲 No 🔿	
Saturation Pres	ent?	Yes \bigcirc No	ullet	Depth (inc	hes):		-	ina nyanology riese		
Describe Reco	orded Data (stre	am daude, mo	onitor we	l, aerial nho	otos. prev	ious insne	ctions). if :	available:		
Remarks:										

Photo Path: C:\Users\dgallagher\Desktop\Wetland_Sample_Point_Photos\



SP1

Plot ID:

Photo File: N e	one.bmp	Orientation:		-facing
Lat/Long or UTM:	Long/Easting: 0		Lat/Northing: 0	
Description:				

WETLAND DETERMINATION DATA FORM - Arid West Region

Project/Site: Guadalupe River Bridge Replacement Project	City/County: San Jose/Santa Clara	County Sampling	Date: 14-Aug-20
Applicant/Owner: Peninsula Corridor Joint Powers Board	State: CA	Sampling Point	t: SP2
Investigator(s): DWG, AB	Section, Township, Range: S	T	R
Landform (hillslope, terrace, etc.): Floodplain	Local relief (concave, convex,	none): concave	Slope: <u>0.0</u> % / <u>0.0</u>
Subregion (LRR): LRR C Lat.:	37.317745 Long.	-121.887926	Datum: WGS84
Soil Map Unit Name: <u>165 – Urbanland-Campbell complex</u> , 0 to 2 percent	nt slopes	NWI classification: P	EM1F
Are climatic/hydrologic conditions on the site typical for this time of ye	ear? Yes $ullet$ No $igodom$ (1	f no, explain in Remarks.)	1
Are Vegetation 🗹 , Soil 🗌 , or Hydrology 🗌 significant	ly disturbed? Are "Normal Ci	rcumstances" present?	Yes 🔾 No 🖲
Are Vegetation 🗌 , Soil 🗌 , or Hydrology 🗌 naturally j	problematic? (If needed, exp	plain any answers in Rema	arks.)

Summary of Findings - Attach site map showing sampling point locations, transects, important features, etc.

Hydrophytic Vegetation Present?	Yes 🖲	No O	Is the Sampled Area	
Hydric Soil Present?	Yes 🖲	No 🔿	15 the Sampled Area	
Wetland Hydrology Present?	Yes 🖲	No 🔿	within a Wetland?	
Remarks: Evidence of Recent Mo	wing			

Dominant

VEGETATION - Use scientific names of plants.

		—Species?			
Tree Stratum (Plot size:	Absolute % Cover	Rel.Strat	Indicator	Dominance Test worksheet:	
			Status	Number of Dominant Species	
1				That are OBL, FACW, or FAC: (A)	
2				Total Number of Dominant	
3	0	0.0%		Species Across All Strata:4_ (B)	
4	0	0.0%			
	0	= Total Co	ver	Percent of dominant Species	
Sapling/Shrub Stratum (Plot size: <u>10 x 10</u>)	-			That Are OBL, FACW, or FAC:	
1. Salix lasiolepis	10	✔ 100.0%	6 FACW	Prevalence Index worksheet:	
2		0.0%		Total % Cover of: Multiply by:	
3.	0	0.0%		OBL species $20 \times 1 = 20$	
4.	0	0.0%		EACW species $20 \times 2 = 40$	
5.	0	0.0%		FAC species $10 \times 3 = 30$	
	10	= Total Co	ver	FACU species $0 \times 4 = 0$	
Herb Stratum (Plot size: 10 x 10)				$0 \times 5 = 0$	
1. Typha latifolia	20	✓ 50.0%	OBL		
2. Mentha spicata	10	25.0%	FACW	$\begin{array}{c} \text{Column Totals:} \underline{50} \text{(A)} \underline{90} \text{(B)} \end{array}$	
3. Artemisia douglasiana	10	✓ 25.0%	FAC	Prevalence Index = B/A = 1.800	
4	0	0.0%		Hydrophytic Vegetation Indicators:	
5.	0	0.0%		\checkmark Dominance Test is > 50%	
6.	0	0.0%		\checkmark Prevalence Index is <3.0 ¹	
7.	0	0.0%			
8.	0	0.0%		data in Remarks or on a separate sheet)	
9.	0	0.0%			
10.	0	0.0%			
11.	0	0.0%			
	40	= Total Co	ver	¹ Indicators of hydric soil and wetland hydrology must	
Woody Vine Stratum (Plot size:)				be present, unless disturbed or problematic.	
1	0	0.0%			
2				Hydrophytic	
		0.0%		Vegetation	
	0	= Total Co	ver	Present? Yes V NO U	
% Bare Ground in Herb Stratum: 0 % Cover of Biotic Crust 0					
Remarks:					

No bare ground, remaining area covered in a layer of thatch (~0.5 inch).

*Indicator suffix = National status or professional decision assigned because Regional status not defined by FWS.

Soil

Sampling Point: SP2

nches)	Color (n	noist)	%	Color	(moist)	_%	Tvpe ¹	Loc ²	Texture	Remarks
)-16	10YR	3/2	90 -	5YR	6/8	10	C		Silty Clay Loam	Soil very moist
			·							
e: C=Con	centration. D=	Depletion	n. RM=Reduc	ed Matrix, Ss. unless	CS=Covere	d or Coate	ed Sand Grai	ins ² Locat	ion: PL=Pore Lining	. M=Matrix
listosol (/	A1)	-ppiloub		Sar	ndy Redox (S5)	,			
Histic Epip	pedon (A2)				ipped Matrix	(S6)				(A10) (LRP B)
Black Histi	ic (A3)				amy Mucky I	Mineral (F	1)			(A10) (ERC B)
Hydrogen	Sulfide (A4)			Loa	amy Gleyed	Matrix (F2	2)		Red Parent	Material (TE2)
Stratified I	Layers (A5) (L	RR C)		🖌 De	pleted Matri	x (F3)			Other (Expl	ain in Remarks)
1 cm Muck	k (A9) (LRR D)			Red	dox Dark Su	rface (F6))		• • • • (_ + •	
Depleted I	Below Dark Su	rface (A1	L)	🗌 De	pleted Dark	Surface (I	F7)			
Thick Dark	< Surface (A12)		Red	dox depress	ions (F8)			3	and a set of a second set of a second
Sandy Muc	ck Mineral (S1)		🗌 Vei	mal Pools (F	9)			wetland hydr	rology must be present.
Sandy Gle	yed Matrix (S4	•)							,	5, 1
trictive La	ayer (if prese	ent):								
Type:		ent):							Hydric Soil Pres	ent? Yes 🖲 No 🔾
Type: Depth (inch narks:	nes):	ent):							Hydric Soil Pres	ent? Yes 🖲 No 🔾
trictive La Type: Depth (inch narks: drology	nes):	ent):							Hydric Soil Pres	ent? Yes • No O
trictive La Type: Depth (inch marks: drology tland Hydr	nes):	ators:							Hydric Soil Pres	ent? Yes • No O
trictive La Type: Depth (inch narks: Irology Iand Hyde Surface W	nes): rology Indica cators (minir ater (A1)	ators:	one required	l; check z	all that app	ly)			Hydric Soil Pres	ent? Yes No O
trictive La Type: Depth (inch narks: Irology Iand Hydu nary India Surface W High Wate	nes): rology Indica cators (minir 'ater (A1) er Table (A2)	ators:	ne requirec	I; check a	all that app alt Crust (B	ly) 11) B12)			Hydric Soil Pres	ent? Yes No Y Indicators (2 or more require Marks (B1) (Riverine) ent Denosits (B2) (Riverine)
rictive La ype: Depth (inch narks: Irology land Hyde nary India Surface W High Wate Saturation	rology Indica cators (minir ater (A1) er Table (A2)	ators:	one required	I; check a	all that app alt Crust (B biotic Crust (aquatic Inve	ly) 11) B12) rtebrates	(B13)		Hydric Soil Press	ent? Yes No ry Indicators (2 or more require Marks (B1) (Riverine) ent Deposits (B2) (Riverine) Deposits (B3) Riverine)
Trictive La ype: Depth (inch narks: Irology land Hyde nary Indic Surface W High Wate Saturation Water Mar	rology Indica cators (minir ater (A1) er Table (A2) (A3) rks (B1) (Nonr	ators: num of c	one requirec	I; check a S B A A H H	all that app Salt Crust (B Siotic Crust (Aquatic Inve Hydrogen Su	ly) 11) B12) rtebrates Ifide Odoo	(B13) r (C1)		Hydric Soil Press	ent? Yes No Y Indicators (2 or more require Marks (B1) (Riverine) ent Deposits (B2) (Riverine) Deposits (B3) Riverine) age Patterns (B10)
Trictive La Type: Depth (inch arks: Trology Iand Hyde nary India Surface W High Wate Saturation Water Mar Sediment	rology Indica cators (minir 'ater (A1) er Table (A2) (A3) rks (B1) (Nonr Deposits (B2)	ators: num of c iverine) (Nonriver	one requirec	I; check a S B A H C C	all that app all that app alt Crust (B Biotic Crust (Aquatic Inve Hydrogen Su Dxidized Rhi	ly) 11) B12) rtebrates Ifide Odor zospheres	(B13) r (C1) : along Livin	 g Roots (C3	Hydric Soil Press	ent? Yes No O
rictive La ype: Depth (inch narks: Irology Iand Hyde nary India Surface W High Wate Saturation Water Mar Sediment Drift depo	rology Indica cators (minir fater (A1) er Table (A2) n (A3) rks (B1) (Nonr Deposits (B2) sits (B3) (Non	ators: num of c (Nonriver eriverine)	one requirec	I; check a S B B B C C C C C C C C C C C C C C C C	all that app alt Crust (B Giotic Crust (Aquatic Inve Hydrogen Su Dxidized Rhi Iresence of	ly) 11) B12) rtebrates lfide Odoi zospheres Reduced I	(B13) r (C1) s along Living ron (C4)	g Roots (C3	Hydric Soil Press	ent? Yes No No ry Indicators (2 or more require Marks (B1) (Riverine) ent Deposits (B2) (Riverine) Deposits (B3) Riverine) age Patterns (B10) eason Water Table (C2) sh Burrows (C8)
trictive La Type: Depth (inch narks: drology :land Hyde nary India Surface W High Wate Saturation Water Mar Sediment Drift depo Surface Sc	rology Indica cators (minir fater (A1) er Table (A2) i (A3) rks (B1) (Nonr Deposits (B2) sits (B3) (Non pil Cracks (B6)	ators: num of c iverine) (Nonriver eriverine)	one requirec	I; check a S B A C P R R	all that app alt Crust (B Siotic Crust (Aquatic Inve Aydrogen Su Aydrogen Su Aydrogen Su Aydrogen Su Aydrogen Su Aydrogen Su Aydrogen Su	ly) 11) B12) rtebrates lfide Odoi zospheres Reduced I Reduction	(B13) r (C1) s along Livin ron (C4) i n Plowed S	g Roots (C3 Soils (C6)	Hydric Soil Press	ent? Yes No No ry Indicators (2 or more require Marks (B1) (Riverine) ent Deposits (B2) (Riverine) Deposits (B3) Riverine) age Patterns (B10) eason Water Table (C2) sh Burrows (C8) ation Visible on Aerial Imagery (C9)
trictive La Type: Depth (inch narks: drology drology darks: drology dro	rology Indica cators (minir ater (A1) er Table (A2) (A3) rks (B1) (Nonr Deposits (B2) sits (B3) (Non pil Cracks (B6) n Visible on Ae	ators: num of c (Nonriver eriverine) erial Image	ine)	l; check a S B B C C C C C C C C C C C C C C C C C	all that app Galt Crust (B Giotic Crust (Aquatic Inve Aydrogen Su Aydrogen Su	ly) 11) B12) rtebrates lfide Odoo zospheres Reduced I Reduction urface (C7	(B13) r (C1) s along Living ron (C4) i n Plowed S	g Roots (C3)	Hydric Soil Press	ent? Yes No Yes No Yey Indicators (2 or more required Marks (B1) (Riverine) ent Deposits (B2) (Riverine) Deposits (B3) Riverine) Deposits (B3) Riverine) Deposits (B10) eason Water Table (C2) sh Burrows (C8) ation Visible on Aerial Imagery (C9) w Aquitard (D3)
trictive La Type: Depth (inch narks: drology dand Hydu nary India Surface W High Wate Saturation Water Mar Sediment Drift depo Surface Sc Inundation Water-Sta	rology Indica cators (minir ater (A1) er Table (A2) (A3) rks (B1) (Nonr Deposits (B2) sits (B3) (Non bil Cracks (B6) n Visible on Ae ined Leaves (E	ators: num of c (Nonriver eriverine) erial Image 39)	one requirec	I; check a S B A C C C C C C C C C C C C C C C C C C	all that app all that app alt Crust (B siotic Crust (Aquatic Inve lydrogen Su lydrogen Su	ly) 11) B12) Ifide Odou zospheres Reduced I Reduction urface (C7 in in Rem	(B13) r (C1) ; along Livin iron (C4) i in Plowed S ') arks)	g Roots (C3 Soils (C6)	Hydric Soil Press	ent? Yes No y Indicators (2 or more require Marks (B1) (Riverine) ent Deposits (B2) (Riverine) Deposits (B3) Riverine) age Patterns (B10) eason Water Table (C2) sh Burrows (C8) ation Visible on Aerial Imagery (C9) w Aquitard (D3) eutral Test (D5)
trictive La Type: Depth (inch marks: drology cland Hyde mary India Surface W High Wate Saturation Water Mar Sediment Drift depo Surface Sc Inundatior Water-Sta d Observa	rology Indica cators (minir dater (A1) er Table (A2) n (A3) rks (B1) (Nonr Deposits (B2) sits (B3) (Non pil Cracks (B6) n Visible on Ae ined Leaves (E	ators: num of c (Nonriver eriverine) erial Imago 39)	ine) ery (B7)	I; check z S B B C C C C C C C C C C C C C C C C C	all that app alt Crust (B Giotic Crust (Aquatic Inve Hydrogen Su Dividized Rhi Dividized Rhi Di Rhi Dividized Rhi Dividized Rhi Dividized Rhi Dividized Rhi	ly) 11) B12) rtebrates lfide Odoi zospheres Reduced I Reduction urface (C7 in in Rem	(B13) r (C1) ; along Livin ; ron (C4) ; in Plowed S ;) arks)	g Roots (C3 Soils (C6)	Hydric Soil Press	ent? Yes No Y Indicators (2 or more require Marks (B1) (Riverine) ent Deposits (B2) (Riverine) Deposits (B3) Riverine) age Patterns (B10) eason Water Table (C2) sh Burrows (C8) tition Visible on Aerial Imagery (C9) w Aquitard (D3) eutral Test (D5)
trictive La Type: Depth (inch marks: drology tland Hyde mary India Surface W High Wate Saturation Water Mar Sediment Drift depo Surface Sc Inundation Water-Sta dd Observa	rology Indica cators (minir 'ater (A1) er Table (A2) (A3) rks (B1) (Nonr Deposits (B2) sits (B3) (Non Dil Cracks (B6) n Visible on Ae ined Leaves (E ations: Present?	ators: num of c iverine) (Nonriver eriverine) erial Imago (89) Yes	ine) ery (B7)		all that app salt Crust (B liotic Crust (liotic Crust (lydrogen Su lydrogen	IV) 11) B12) rtebrates Ifide Odou zospheres Reduced I Reduction urface (C7 in in Rem. hes):	(B13) r (C1) s along Living ron (C4) i n Plowed S) arks)	g Roots (C3)	Hydric Soil Press	ent? Yes No No ry Indicators (2 or more require Marks (B1) (Riverine) ent Deposits (B2) (Riverine) Deposits (B3) Riverine) age Patterns (B10) eason Water Table (C2) sh Burrows (C8) ation Visible on Aerial Imagery (C9) w Aquitard (D3) eutral Test (D5)
trictive La Type: Depth (inch marks: drology tland Hyde mary India Surface W High Wate Saturation Water Mar Sediment Drift depo Surface Sc Inundation Water-Sta dobserva face Water	rology Indica cators (minir ater (A1) er Table (A2) (A3) rks (B1) (Nonr Deposits (B2) sits (B3) (Non bil Cracks (B6) n Visible on Ae ined Leaves (E ations: Present?	ators: num of c (Nonriver eriverine) erial Imagu 39) Yes (Yes (one requirec ine) ery (B7) No (e) No (e)		all that app all that app alt Crust (B siotic Crust (Aquatic Inve lydrogen Su dydrogen Su	ly) 11) B12) rtebrates - Ifide Odor zospheres Reduced I Reduction urface (C7 in in Rem- nes): nes):	(B13) r (C1) s along Living ron (C4) i n Plowed S) arks)	g Roots (C3) Soils (C6)	Hydric Soil Press	ent? Yes No
trictive La Type: Depth (inch marks: 	rology Indica cators (minir dater (A1) er Table (A2) (A3) rks (B1) (Nonr Deposits (B2) sits (B3) (Non bil Cracks (B6) n Visible on Ae ined Leaves (E ations: Present? resent? resent? ary fringe)	ators: num of c (Nonriver eriverine) erial Imago 39) Yes (Yes (Yes (ine) ery (B7) No @ No @ No @	I; check z S B B C C C C C C C C C C C C C C C C C	all that app alt Crust (B Giotic Crust (Aquatic Inve Aydrogen Su Dxidized Rhi Presence of Agreent Iron Thin Muck Su Dther (Expla Depth (incl Depth (incl Depth (incl	IV) 11) B12) rtebrates Ifide Odou zospheres Reduced I Reduction urface (C7 in in Remu- nes): nes): nes):	(B13) r (C1) s along Livin ron (C4) in Plowed S ') arks)	g Roots (C3 Soils (C6)	Hydric Soil Press	ent? Yes No ry Indicators (2 or more require Marks (B1) (Riverine) ent Deposits (B2) (Riverine) Deposits (B3) Riverine) age Patterns (B10) eason Water Table (C2) sh Burrows (C8) tion Visible on Aerial Imagery (C9) w Aquitard (D3) eutral Test (D5) sent? Yes No

 Pot Dir
 SP2
 Photo Path: C:\Users\dgallagher\Desktop\Wetland_Sample_Point_Photos\

Lat/Long or UTM : Long/Easting: -121.887926 Lat/Northing: 37.317745
Description:

No Photo

Photo File: No	one.bmp	Orientation:		-facing
Lat/Long or UTM:	Long/Easting: 0		Lat/Northing: 0	
Description:				

WETLAND DETERMINATION DATA FORM - Arid West Region

Project/Site: Guadalupe River Bridge Replacement Project	City/County: San Jose/Santa Clar	a County Sampling	Sampling Date: <u>14-Aug-20</u>	
Applicant/Owner: Peninsula Corridor Joint Powers Board	State: CA	Sampling Poin	ıt: SI	P3
Investigator(s): DWG, AB	Section, Township, Range: S	;т	R	
Landform (hillslope, terrace, etc.): Floodplain	Local relief (concave, convex,	none): concave	Slope: 0.(<u>)</u> %/°
Subregion (LRR): LRR C Lat.:	37.319288 Long	: -121.888767	Datum: W	GS84
Soil Map Unit Name: <u>165 – Urbanland-Campbell complex, 0 to 2 perce</u>	nt slopes	NWI classification: P	PEM1F	
Are climatic/hydrologic conditions on the site typical for this time of ye	ear? Yes \odot No \bigcirc (If no, explain in Remarks.)) Vac 🔍 Na	\bigcirc
Are Vegetation , Soil , or Hydrology significant	ay disturbed? Are "Normal (ircumstances" present?	Tes 🔄 NU	0
	problematic: (If needed, e)	plain any answers in Rem	arks.)	

Summary of Findings - Attach site map showing sampling point locations, transects, important features, etc.

Hydrophytic Vegetation Present?	Yes 🖲	No 🔿	Is the Sampled Area	
Hydric Soil Present?	Yes 🖲	No 🔿	within a Watland2	Yes 🔍 No 🔿
Wetland Hydrology Present?	Yes 🖲	No \bigcirc	within a wetland?	
Remarks:				

(Plot size:		Rel.Strat.	Indicator	Dominance Test worksheet:
	% Cover		Status	Number of Dominant Species
	0			That are OBL, FACW, or FAC: (A)
	0	0.0%		Total Number of Dominant
·	0	0.0%		Species Across All Strata: (B)
	0	0.0%		
pling/Shrub Stratum (Plot size:)	0	= Total Cov	er	Percent of dominant Species That Are OBL, FACW, or FAC: <u>100.0%</u> (A/B
· · · · · · · · · · · · · · · · · · ·	0	0.0%		Prevalence Index worksheet:
	0	0.0%		Total % Cover of: Multiply by:
	0	0.0%		$\frac{1000}{1000} = \frac{1000}{1000} = \frac{1000}{1000$
	0			$\frac{1}{1000} = \frac{1}{1000} = \frac{1}{1000} = \frac{1}{10000} = \frac{1}{10000000000000000000000000000000000$
	0			FACW species $0 \times 2 = 0$
			~~	FAC species $10 \times 3 = 30$
b Stratum (Plot size: <u>5 x 5</u>)	0		er	FACU species $0 \times 4 = 0$
Ludwigia peploides	60	✔ 84.5%	OBL	UPL species $x 5 = 3$
. Xanthium strumarium	5	7.0%	FAC	Column Totals: <u>71</u> (A) <u>95</u> (B
. Lotus corniculatus	5	7.0%	FAC	Prevalence Index = $B/A = 1.338$
. Melilotus alba	1	1.4%	UPL	Hydrophytic Vegetation Indicators:
	0	0.0%		\checkmark Dominance Test is > 50%
	0	0.0%		\checkmark Prevalence Index is <3.0 ¹
	0	0.0%		
	0	0.0%		data in Remarks or on a separate sheet)
	0	0.0%		Droblomatic Hydrophytic Vegetation ¹ (Explain)
	0	0.0%		
	0	0.0%		
	71	= Total Cov	er	¹ Indicators of hydric soil and wetland hydrology must be present, unless disturbed or problematic.
oodv Vine Stratum (Plot size:)		_		
	0	0.0%		
·	0	0.0%		Hydrophytic
	0	= Total Cov	er	Present? Yes I No
% Bare Ground in Herb Stratum: 29 %	Cover of Bioti	ic Crust ∩		

*Indicator suffix = National status or professional decision assigned because Regional status not defined by FWS.

Soil

Sampling Point: SP3

Profile Description: (Describe to the dep	oth needed to document	the indicator or co	nfirm the a	bsence of indicators.)	
Depth Matrix	Rec	ox Features			
(inches) Color (moist) %	Color (moist)	<u>% Tvpe¹</u>	Loc ²	Texture	Remarks
0-16 10YR 5/1 90	5YR5/8	10 C	PL	Silty Clay	
· · ·					
				-	
¹ Type: C=Concentration. D=Depletion. RM=1	Reduced Matrix, CS=Covere	d or Coated Sand Gra	ns ² Locatio	on: PL=Pore Lining. M=Matri	x
Hydric Soil Indicators: (Applicable to a	II LRRs, unless otherwis	e noted.)		Indicators for Problem	atic Hydric Soils: ³
Histosol (A1)	Sandy Redox (S5)		1 cm Muck (A9) (LRR	C)
Histic Epipedon (A2)	Stripped Matrix	< (S6)		2 cm Muck (A10) (LRI	Я В)
Black Histic (A3)	Loamy Mucky	Mineral (F1)		Reduced Vertic (F18)	
	Loamy Gleyed	Matrix (F2)		Red Parent Material (TF2)
	Depleted Matri	x (F3)		Other (Explain in Rem	narks)
	Redox Dark Su	rface (F6)			
	Depleted Dark	Surface (F7)			
	Redox depress	ions (F8)		3	
Sandy Muck Mineral (S1)	Vernal Pools (I	-9)		 Indicators of hydrophyti wetland hydrology mu 	c vegetation and st be present.
Sandy Gleyed Matrix (S4)				nedana nyarology ma	
Restrictive Layer (if present):					
Туре:					
Depth (inches):				Hydric Soil Present?	Yes 🔍 No 🔾
Remarks:					
Soil very moist likley evidence of recent	inundation				
Hydrology					
пушоюду					
Wetland Hydrology Indicators:					
Primary Indicators (minimum of one red	uired; check all that app	oly)		Secondary Indicat	cors (2 or more required)
Surface Water (A1)	Salt Crust (B	11)		Water Marks (B	1) (Riverine)
High Water Table (A2)	Biotic Crust	(B12)		Sediment Depos	sits (B2) (Riverine)
Saturation (A3)	Aquatic Inve	rtebrates (B13)		Drift Deposits (I	33) Riverine)
Water Marks (B1) (Nonriverine)	Hvdrogen Su	lfide Odor (C1)		Drainage Patter	ns (B10)
Sediment Deposits (B2) (Nonriverine)	Oxidized Rhi	zospheres along Livin	a Roots (C3)	Dry Season Wat	er Table (C2)
$\square \text{ Drift deposits (B3) (Noneriverine)}$		Reduced Iron (C4)	,	Cravfish Burrow	e(2)
Surface Soil Cracks (B6)		Reduction in Plowed 9	oils (C6)	Saturation Visib	le on Aerial Imageny (CQ)
 Jourdation Visible on Aerial Imagen/ (B7) Thin Muck S	urface (C7)			
Water-Stained Leaves (B9)) Cther (Evola	in in Pemarke)			t (DS)
Field Observations:					
Surface Water Present? Yes \cup	Depth (incl	nes):	_		
Water Table Present? Yes O	lo 💿 Depth (incl	nes):	_		
Saturation Present? Yes O	lo Depth (incl	nes):	Wetlan	nd Hydrology Present?	Yes 🔍 No 🔾

Remarks:

Describe Recorded Data (stream gauge, monitor well, aerial photos, previous inspections), if available:

 Pot ID:
 \$P3
 Photo Path: C:\Users\dgallagher\Desktop\Wetland_Sample_Point_Photos\

Photo File: SI	P3.JPG	Orientation:	-facing
Lat/Long or UTM :	Long/Easting: -1	21.888767	Lat/Northing: 37.319288
Description:			

one.bmp	Orientation:		-facing
Long/Easting: 0		Lat/Northing: 0	
	one.bmp Long/Easting: 0	Done.bmp Orientation: Long/Easting: 0	Done.bmp Orientation: Long/Easting: 0 Lat/Northing: 0

WETLAND DETERMINATION DATA FORM - Arid West Region

Project/Site: Guadalupe River Bridge Replacement Project	City/County: San Jose/Santa Clara	County Sampling	Date: 14-Aug-20
Applicant/Owner: Peninsula Corridor Joint Powers Board	State: CA	Sampling Point	t: SP4
Investigator(s): DWG, AB	Section, Township, Range: S	T	R
Landform (hillslope, terrace, etc.): Floodplain	Local relief (concave, convex,	none): convex	Slope: <u>30.0</u> % / <u>16.7</u> °
Subregion (LRR): LRR C Lat.:	37.31933 Long.:	-121.888632	Datum: WGS84
Soil Map Unit Name: <u>165 – Urbanland-Campbell complex</u> , 0 to 2 percent	nt slopes	NWI classification:	
Are climatic/hydrologic conditions on the site typical for this time of ye	ear? Yes $ullet$ No $igodom$ (I	f no, explain in Remarks.)	1
Are Vegetation 🗹 , Soil 🗌 , or Hydrology 🗌 significant	tly disturbed? Are "Normal Ci	rcumstances" present?	Yes 🔾 No 🖲
Are Vegetation 🗌 , Soil 🗌 , or Hydrology 🗌 naturally j	problematic? (If needed, exp	plain any answers in Rema	arks.)

Summary of Findings - Attach site map showing sampling point locations, transects, important features, etc.

Hydrophytic Vegetation Present?	Yes \bigcirc	No 🖲	Is the Sampled Area	
Hydric Soil Present?	Yes 🖲	No 🔿	within a Watland2	Yes 🔿 No 🔍
Wetland Hydrology Present?	Yes \bigcirc	No 🖲	within a wetland?	
Remarks: Evidence of Recent Mov	wing			

Dominant

VEGETATION - Use scientific names of plants.

Trop Stratum (Plot size)	Absolute	Rel.Strat.	Indicator	Dominance Test worksheet:
1	<u>-70 COVEI</u>		Status	Number of Dominant Species
2		0.0%		$\frac{1}{1}$
2				Total Number of Dominant
<i>A</i>				Species Across All Strata: (B)
т	0	0.0%		Percent of dominant Species
Sanling/Shrub Stratum (Plot size:)	0	= Total Cove	r	That Are OBL, FACW, or FAC: 0.0% (A/B)
1	0	0.0%		Provalence Index worksheet:
2.	0	0.0%		Total % Cover of Multiply by:
3.	0	0.0%		$\begin{array}{c} 10 \text{ cover or} \\ \hline \end{array} \\ \hline \\ \\ \hline \end{array} \\ \\ \hline \\ \\ \hline \end{array} \\ \\ \hline \\ \\ \hline \\ \\ \hline \\ \\ \hline \end{array} \\ \\ \hline \\ \\ \hline \\ \\ \hline \\ \\ \hline \\ \\ \\ \\$
4.				$\frac{\partial \partial E}{\partial x} = \frac{\partial E}{\partial x} = \frac{\partial E}{\partial x}$
5.				FACW spectres 0 x 2 = 0
		Cours		FAC species $0 \times 3 = 0$
Herb Stratum (Plot size: 5 x 5)	0		۲ .	FACU species $____$
1 Convolvulus arvensis	50	✔ 100.0%	UPL	UPL species -50 x 5 = -250
2	0	0.0%		Column Totals: <u>50</u> (A) <u>250</u> (B)
3.	0	0.0%		Prevalence Index = $B/A = 5.000$
4.	0	0.0%		Hydrophytic Vegetation Indicators:
5.	0	0.0%		Dominance Test is > 50%
6.	0	0.0%		$\square Prevalence Index is <3.0^{1}$
7	0	0.0%		
8	0	0.0%		data in Remarks or on a separate sheet)
9.	0	0.0%		Problematic Hydrophytic Vegetation ¹ (Explain)
10.	0	0.0%		
11	0	0.0%		
	50	= Total Cove	r	¹ Indicators of hydric soil and wetland hydrology must be present, unless disturbed or problematic.
Woodv Vine Stratum (Plot size:)				
1	0	0.0%		
2	0	0.0%		Hydrophytic
	0	= Total Cove	r	Present? Yes No 🔍
% Bare Ground in Herb Stratum: _0 % Co	ver of Bioti	ic Crust 0		
Remarks:				

No bare ground, remaining area covered in a layer of thatch (~ 1 inch).

*Indicator suffix = National status or professional decision assigned because Regional status not defined by FWS.

Soil

Sampling Point: SP4

Profile Descri	ption: (Describe to	the depth nee	ded to document	the indica	tor or confi	irm the	absence of indicators.)	
Depth Matrix Redox Features								
(inches)	Color (moist)	%	Color (moist)	%	Tvpe ¹	Loc ²	Texture	Remarks
0-16	10YR 3/2	90	5YR 4/6	10	D	М	Silty Clay	Relic redox
	<u></u>							
¹ Type: C=Conc	entration. D=Depletic	on. RM=Reduced	I Matrix, CS=Covered	d or Coated	Sand Grains	² Locat	tion: PL=Pore Lining. M=	Matrix
Hydric Soil In	ndicators: (Applica	ble to all LRRs	, unless otherwise	e noted.)			Indicators for Prob	lematic Hydric Soils: ³
Histosol (A	1)		Sandy Redox (S5)			1 cm Muck (A9)	(LRR C)
Histic Epipe	edon (A2)		Stripped Matrix	(S6)			2 cm Muck (A10)	(LRR B)
Black Histic	c (A3)		Loamy Mucky I	Mineral (F1))		Reduced Vertic (F18)
			Loamy Gleyed	Matrix (F2)			Red Parent Mate	rial (TF2)
	ayeis (AD) (LKK C)		Depleted Matri	x (F3)			Other (Explain in	Remarks)
	(MJ) (LRR D)	11)	Redox Dark Su	rface (F6)				
	Surface (A12)	"	Depleted Dark	Surface (F7	7)			
Sandy Muc	k Mineral (S1)		Redox depress	ions (F8)			³ Indicators of hydror	phytic vegetation and
Sandy Glev	red Matrix (S4)		Vernal Pools (F	9)			wetland hydrology	y must be present.
Restrictive La	ver (if present):							
Type ¹	yer (ii present).							
Depth (inch	ec).						Hydric Soil Present?	Yes 💿 No 🔾
Pomarke:).							
Dolio rodov lik	by from oversition	of ovicitor floo	delain and donasi	tod coile t	a form the	hanko		
Relic Teuox-like						Dariks.		
Hydrology	,							
Wetland Hydr	ology Indicators:							
Primary Indic	ators (minimum of	one required:	check all that app	lv)			Secondary In	dicators (2 or more required)
Surface Wa	ater (A1)		Salt Crust (B	11)			Water Mark	(B1) (Riverine)
🗌 High Water	r Table (A2)		🗌 Biotic Crust (B12)			Sediment D	Deposits (B2) (Riverine)
Saturation	(A3)		Aquatic Inve	rtebrates (E	813)		Drift Depos	sits (B3) Riverine)
Water Mark	ks (B1) (Nonriverine)		🗌 Hydrogen Su	lfide Odor ((C1)		Drainage P	atterns (B10)
Sediment D	Deposits (B2) (Nonrive	erine)	Oxidized Rhi	zospheres a	along Living R	Roots (C3	B) Dry Season	Water Table (C2)
Drift depos	its (B3) (Noneriverine)	Presence of I	Reduced Irc	on (C4)		Crayfish Bu	rrows (C8)
Surface Soi	il Cracks (B6)		Recent Iron I	Reduction i	n Plowed Soil	ls (C6)	Saturation	Visible on Aerial Imagery (C9)
Inundation	Visible on Aerial Imag	gery (B7)	Thin Muck Su	urface (C7)			Shallow Aq	uitard (D3)
Water-Stain	ned Leaves (B9)		Other (Expla	in in Remar	ks)		FAC-neutra	l Test (D5)
Field Observa	tions:							
Surface Water P	Present? Yes	○ No	Depth (incl	nes):				
Water Table Pre	esent? Yes	O No 🖲	Donth (incl					
Saturation Prese	ent?					Wetla	and Hydrology Present	? Yes 🔾 No 🖲
(includes capilla	ary fringe) Yes		Depth (incl	nes):				
Describe Reco	orded Data (stream	gauge, monito	or well, aerial pho	tos, previo	ous inspecti	ons), if	available:	
Remarks:								

Plot ID:	SP4

Photo Path: C:\Users\dgallagher\Desktop\Wetland_Sample_Point_Photos\



Photo File: 51	4.JPG	Unentation.	-racing
Lat/Long or UTM :	Long/Easting	: -121.888632	Lat/Northing: 37.31933
Description:			

No Phot	D
---------	---

Photo File: No	one.bmp	Orientation:		-facing
Lat/Long or UTM:	Long/Easting: 0		Lat/Northing: 0	
Description:				
l				

Appendix D: Photographic Documentation of the Study Area



Photo 1. Looking downstream at Guadalupe MT-2 track bridge. Guadalupe River and associated riparian habitat are clearly visible.



Photo 2. Looking south at the flood control basin. August 2020.



Photo 3. Seasonal wetland within the flood control basin. August 2020.



Photo 4. Perennial Wetland within the flood control basin. A large culvert empties into the basin from under Edwards Avenue at McClellan Avenue. August 2020.



Photo 5. Flood control basin with annual grassland habitat along the banks. The area was recently mowed as indicated by the remaining thatch. The green vegetation in the bottom of the basin is a seasonal wetland. August 2020.



Photo 6. Gravel access road for flood control basin. This area was mapped as developed habitat.



Photo 7. Guadalupe River. Aquatic and riparian habitat are clearly visible.

Appendix E: Aquatic Resources Table
Waters Name	State	Cowardin Code	HGM Code	Measurement Type	Amount	Units	Water Type	Latitude	Longitude	Local Waterway
PS1	CALIFORNIA	R2UB	RIVERINE	Area	1.45	ACRE	TNW	37.316792°	-121.888145°	Guadalupe River
PM1	CALIFORNIA	PEM1F	DEPRESS	Area	1.61	ACRE	TNWW	37.317860°	-121.888278°	Guadalupe River
SW1	CALIFORNIA	PEM1E	DEPRESS	Area	0.35	ACRE	TNWW	37.317248°	-121.887835°	Guadalupe River
SW2	CALIFORNIA	PEM1E	DEPRESS	Area	0.98	ACRE	TNWW	37.318715°	-121.888443°	Guadalupe River